Rabbit anti-Human Tumor necrosis factor receptor superfamily member 10D Polyclonal Antibody | anti-TNFRSF10D antibody

Rabbit anti-human Tumor necrosis factor receptor superfamily member 10D polyclonal Antibody(TNFRSF10D)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA119150	**Rabbit**	**Human**	**Immunohistochemistry, Western Blot, ELISA**

PRODUCT SYNONYMS

Tumor necrosis factor receptor superfamily member 10D; Polyclonal Antibody; Rabbit anti-human Tumor necrosis factor receptor superfamily member 10D polyclonal Antibody(TNFRSF10D); Decoy receptor 2; DcR2; TNF-related apoptosis-inducing ligand receptor 4; TRAIL receptor 4; TRAIL-R4; TRAIL receptor with a truncated death domain; CD264; TNFRSF10D; DCR2; TRAILR4; TRUNDD; UNQ251/PRO288; anti-TNFRSF10D antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Sequence Length	386
Applicable Applications for anti-TNFRSF10D antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Conjugate	Non-conjugated
Immunogen	Recombinant human Tumor necrosis factor receptor superfamily member 10D protein (257-386aa)
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-11638 / sc-7550 / sc-7551 / sc-26535 / sc-26534 / sc-28985
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES

IHC (Immunohistochemisry)
Immunohistochemistry of paraffin-embedded human kidney using AAA119150 at dilution 1:100

IHC (Immunohiostchemistry)
Immunohistochemistry of paraffin-embedded human tonsil using AAA119150 at dilution 1:100



WB (Western Blot)
All lanes: Tumor necrosis factor receptor superfamily member 10D antibody at 2ug/ml+mouse liver tissueSecondaryGoat polyclonal to rabbit at 1/10000 dilutionPredicted band size: 42kDaObserved band size: 42kDa

ADDITIONAL INFORMATION

Related Product Information for anti-TNFRSF10D antibody	Receptor for the cytotoxic ligand TRAIL. Contains a truncated death domain and hence is not capable of inducing apoptosis but protects against TRAIL-mediated apoptosis. Reports are contradictory with regards to its ability to induce the NF-kappa-B pathway. According to PubMed:9382840, it cannot but according to PubMed:9430226, it can induce the NF-kappa-B pathway.

NCBI AND UNIPROT INFORMATION

NCBI GI #	22547108
NCBI GeneID	8793
NCBI Accession #	NP_003831.2
NCBI GenBank Nucleotide #	NM_003840.4
UniProt Accession #	Q9UBN6; Q9Y6Q4; B2R8W0
Molecular Weight	41,823 Da
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 10D
NCBI Official Synonym Full Names	tumor necrosis factor receptor superfamily member 10d
NCBI Official Symbol	TNFRSF10D
NCBI Official Synonym Symbols	DCR2; CD264; TRUNDD; TRAILR4; TRAIL-R4
NCBI Protein Information	tumor necrosis factor receptor superfamily member 10D
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 10D
UniProt Gene Name	**TNFRSF10D**
UniProt Synonym Gene Names	DCR2; TRAILR4; TRUNDD; DcR2; TRAIL receptor 4; TRAIL-R4
UniProt Entry Name	TR10D_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: https://stage.aaabiotech.com/polyclonal/tumor-necrosis-factor-receptor-superfamily-member-10d-antibody-immunohistochemistry-western-blot-elisa/119150