

Rabbit anti-Human Rapamycin-insensitive companion of mTOR Polyclonal Antibody | anti-RICTOR antibody

Rabbit anti-human Rapamycin-insensitive companion of mTOR polyclonal Antibody(RICTOR)

CATALOG # AAA119430	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

Rapamycin-insensitive companion of mTOR; Polyclonal Antibody; Rabbit anti-human Rapamycin-insensitive companion of mTOR polyclonal Antibody(RICTOR); AVO3 homolog; hAVO3; RICTOR; anti-RICTOR antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Sequence Length	1732
Applicable Applications for anti-RICTOR antibody	IHC (Immunohistochemistry), ELISA
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Conjugate	Non-conjugated
Immunogen	Recombinant human Rapamycin-insensitive companion of mTOR protein(1-245aa)
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-50678 / sc-130865 / sc-99004
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human spleen using AAA119430 at dilution 1:100



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human lung cancer using AAA119430 at dilution 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-RICTOR antibody	Subunit of mTORC2, which regulates cell growth and survival in response to hormonal signals. mTORC2 is activated by growth factors, but, in contrast to mTORC1, seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. mTORC2 plays a critical role in AKT1 'Ser-473' phosphorylation, which may facilitate the phosphorylation of the activation loop of AKT1 on 'Thr-308' by PDK1 which is a prerequisite for full activation. mTORC2 regulates the phosphorylation of SGK1 at 'Ser-422'. mTORC2 also modulates the phosphorylation of PRKCA on 'Ser-657'. Plays an essential role in embryonic growth and development.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	550544217
NCBI GeneID	253260
NCBI Accession #	NP_001272368.1
NCBI GenBank Nucleotide #	NM_001285439.1
UniProt Accession #	Q6R327; Q68DT5; Q86UB7; Q8N3A0; Q8NCM6; B2RNX0; B7ZMF7
Molecular Weight	195,011 Da
NCBI Official Full Name	rapamycin-insensitive companion of mTOR isoform 2
NCBI Official Synonym Full Names	RPTOR independent companion of MTOR, complex 2
NCBI Official Symbol	RICTOR
NCBI Official Synonym Symbols	PIA; AVO3; hAVO3
NCBI Protein Information	rapamycin-insensitive companion of mTOR
UniProt Protein Name	Rapamycin-insensitive companion of mTOR
UniProt Gene Name	RICTOR
UniProt Synonym Gene Names	hAVO3
UniProt Entry Name	RICTR_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/rapamycin-insensitive-companion-of-mtor-antibody-immunohistochemistry-elisa/119430>