

Rabbit anti-Human RIPK2 Polyclonal Antibody | anti-RIPK2 antibody

RIPK2 Antibody

CATALOG # AAA119528	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

RIPK2; Polyclonal Antibody; RIPK2 Antibody; Receptor-interacting serine/threonine-protein kinase 2; CARD-containing interleukin-1 beta-converting enzyme-associated kinase; CARD-containing IL-1 beta ICE-kinase; RIP-like-interacting CLARP kinase; Receptor-interacting protein 2; RIP-2; Tyrosine-protein kinase RIPK2; CARDIAK; RICK; RIP2; anti-RIPK2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Sequence Length	540
Applicable Applications for anti-RIPK2 antibody	IHC (Immunohistochemistry), ELISA
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Conjugate	Non-conjugated
Immunogen	Recombinant human Receptor-interacting serine/threonine-protein kinase 2 protein (1-540aa)
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-8611 / sc-8610 / sc-22763
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human placenta using AAA119528 at dilution 1:100



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human prostate using AAA119528 at dilution 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-RIPK2 antibody	Serine/threonine/tyrosine kinase that plays an essential role in modulation of innate and adaptive immune responses. Upon stimulation by bacterial peptidoglycans, NOD1 and NOD2 are activated, oligomerize and recruit RIPK2 through CARD-CARD domains. Contributes to the tyrosine phosphorylation of the guanine exchange factor ARHGEF2 through Src tyrosine kinase leading to NF-kappaB activation by NOD2. Once recruited, RIPK2 autophosphorylates and undergoes 'Lys-63'-linked polyubiquitination by E3 ubiquitin ligases XIAP, BIRC2 and BIRC3. The polyubiquitinated protein mediates the recruitment of MAP3K7/TAK1 to IKBKG/NEMO and induces 'Lys-63'-linked polyubiquitination of IKBKG/NEMO and subsequent activation of IKBKB/IKKB. In turn, NF-kappa-B is released from NF-kappa-B inhibitors and translocates into the nucleus where it activates the transcription of hundreds of genes involved in immune response, growth control, or protection against apoptosis. Plays also a role during engagement of the T-cell receptor (TCR) in promoting BCL10 phosphorylation and subsequent NF-kappa-B activation.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506537
NCBI GeneID	8767
NCBI Accession #	NP_003812.1
NCBI GenBank Nucleotide #	NM_003821.5
UniProt Accession #	O43353; Q6UWF0; B7Z748
Molecular Weight	45,582 Da
NCBI Official Full Name	receptor-interacting serine/threonine-protein kinase 2
NCBI Official Synonym Full Names	receptor interacting serine/threonine kinase 2
NCBI Official Symbol	RIPK2
NCBI Official Synonym Symbols	CCK; RICK; RIP2; CARD3; GIG30; CARDIAK
NCBI Protein Information	receptor-interacting serine/threonine-protein kinase 2
UniProt Protein Name	Receptor-interacting serine/threonine-protein kinase 2
UniProt Gene Name	RIPK2
UniProt Synonym Gene Names	CARDIAK; RICK; RIP2; CARD-containing IL-1 beta ICE-kinase; RIP-2
UniProt Entry Name	RIPK2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ripk2-antibody-immunohistochemistry-elisa/119528>