

Rabbit anti-Human BDNF/NT-3 growth factors receptor Polyclonal Antibody | anti-NTRK2 antibody

Rabbit anti-human BDNF/NT-3 growth factors receptor polyclonal Antibody(NTRK2)

CATALOG # AAA119632	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

BDNF/NT-3 growth factors receptor; Polyclonal Antibody; Rabbit anti-human BDNF/NT-3 growth factors receptor polyclonal Antibody(NTRK2); GP145-TrkB; Trk-B; Neurotrophic tyrosine kinase receptor type 2; TrkB tyrosine kinase; Tropomyosin-related kinase B; NTRK2; TRKB; anti-NTRK2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Sequence Length	477
Applicable Applications for anti-NTRK2 antibody	IHC (Immunohistochemistry), ELISA
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Conjugate	Non-conjugated
Immunogen	Recombinant human BDNF/NT-3 growth factors receptor protein(32-282aa)
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-20542 / sc-12 / sc-119 / sc-8316
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human kidney using AAA119632 at dilution 1:100



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human breast cancer using AAA119632 at dilution 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-NTRK2 antibody	Receptor tyrosine kinase involved in the development and the maturation of the central and the peripheral nervous systems through regulation of neuron survival, proliferation, migration, differentiation, and synapse formation and plasticity. Receptor for BDNF/brain-derived neurotrophic factor and NTF4/neurotrophin-4. Alternatively can also bind NTF3/neurotrophin-3 which is less efficient in activating the receptor but regulates neuron survival through NTRK2. Upon ligand-binding, undergoes homodimerization, autophosphorylation and activation. Recruits, phosphorylates and/or activates several downstream effectors including SHC1, FRS2, SH2B1, SH2B2 and PLCG1 that regulate distinct overlapping signaling cascades. Through SHC1, FRS2, SH2B1, SH2B2 activates the GRB2-Ras-MAPK cascade that regulates for instance neuronal differentiation including neurite outgrowth. Through the same effectors controls the Ras-PI3 kinase-AKT1 signaling cascade that mainly regulates growth and survival. Through PLCG1 and the downstream protein kinase C-regulated pathways controls synaptic plasticity. Thereby, plays a role in learning and memory by regulating both short term synaptic function and long-term potentiation. PLCG1 also leads to NF-Kappa-B activation and the transcription of genes involved in cell survival. Hence, it is able to suppress anoikis, the apoptosis resulting from loss of cell-matrix interactions. May also play a role in neurotrophin-dependent calcium signaling in glial cells and mediate communication between neurons and glia.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	55956790
NCBI GeneID	4915
NCBI Accession #	NP_001007098.1
NCBI GenBank Nucleotide #	NM_001007097.2
UniProt Accession #	Q16620; Q16675; Q59GJ1; Q8WXJ5; Q8WXJ6; Q8WXJ7; B1ANZ4; B4DFV9
Molecular Weight	35,332 Da
NCBI Official Full Name	BDNF/NT-3 growth factors receptor isoform b
NCBI Official Synonym Full Names	neurotrophic tyrosine kinase, receptor, type 2
NCBI Official Symbol	NTRK2
NCBI Official Synonym Symbols	TRKB; trk-B; GP145-TrkB
NCBI Protein Information	BDNF/NT-3 growth factors receptor
UniProt Protein Name	BDNF/NT-3 growth factors receptor
UniProt Gene Name	NTRK2
UniProt Synonym Gene Names	TRKB; Trk-B
UniProt Entry Name	NTRK2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.