

Porcine Procollagen Type I Intact N-terminal Propeptide (P1NP) ELISA Kit | P1NP elisa kit

Porcine Procollagen Type I Intact N-terminal Propeptide (P1NP) ELISA Kit

CATALOG #
AAA121358

REACTIVITY
Porcine


PRODUCT SYNONYMS

Procollagen Type I Intact N-terminal Propeptide (P1NP); Porcine Procollagen Type I Intact N-terminal Propeptide (P1NP) ELISA Kit; P1NP elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Porcine
Samples	Serum, plasma, cell culture supernates, cell lysates, tissue homogenates
Assay Type	Quantitative Sandwich
Detection Range	0.5ng/ml - 200ng/ml
Sensitivity	0.28ng/ml
Intra-assay Precision	Intra-Assay Precision (Precision within an assay) Three samples of known concentration were tested on one plate to assess intra-assay precision. Intra-Assay: CV<8%
Inter-assay Precision	Inter-Assay Precision (Precision between assays) Three samples of known concentration were tested in separate assays to assess inter-assay precision. CV(%) = SD/mean x 100. Inter-Assay: CV<10%
Preparation and Storage	Store at 2 to 8 degree C for 6 months.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for P1NP elisa kit

Intended Uses: This sandwich kit is for the accurate quantitative detection of Porcine Procollagen Type I Intact N-terminal Propeptide (also known as P1NP) in serum, plasma, cell culture supernates, cell lysates, tissue homogenates.

Principle of the Assay: This kit is an Enzyme-Linked Immunosorbent Assay (ELISA). The plate has been pre-coated with Porcine P1NP antibody. P1NP present in the sample is added and binds to antibodies coated on the wells. And then biotinylated Porcine P1NP Antibody is added and binds to P1NP in the sample. Then Streptavidin-HRP is added and binds to the Biotinylated P1NP antibody. After incubation unbound Streptavidin-HRP is washed away during a washing step. Substrate solution is then added and color develops in proportion to the amount of Porcine P1NP. The reaction is terminated by addition of acidic stop solution and absorbance is measured at 450 nm.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.