

## Rat Mast Cell Tryptase 4 (MCT4) ELISA Kit | MCT4 elisa kit

Rat Mast Cell Tryptase 4 (MCT4) ELISA Kit

CATALOG #  
**AAA122561**

REACTIVITY  
**Rat**


### PRODUCT SYNONYMS

Mast Cell Tryptase 4 (MCT4); Rat Mast Cell Tryptase 4 (MCT4) ELISA Kit; MCT4 elisa kit

### PRODUCT SPECIFICATIONS

|                         |                                                                          |
|-------------------------|--------------------------------------------------------------------------|
| Reactivity              | Rat                                                                      |
| Samples                 | Serum, plasma, cell culture supernates, cell lysates, tissue homogenates |
| Assay Type              | Quantitative Sandwich                                                    |
| Detection Range         | 0.1ng/ml - 40ng/ml                                                       |
| Sensitivity             | 0.052ng/ml                                                               |
| Intra-Assay             | CV<8%                                                                    |
| Inter-Assay             | CV<10%                                                                   |
| Preparation and Storage | Store at 2 to 8 degree C for 6 months.                                   |

### IMAGES



Standard Curve (Sample)

### ADDITIONAL INFORMATION

#### Related Product Information for MCT4 elisa kit

**Intended Uses:** This sandwich kit is for the accurate quantitative detection of Rat Mast Cell Tryptase 4 (also known as MCT4) in serum, plasma, cell culture supernates, cell lysates, tissue homogenates.

**Principles of the Assay:** This kit is an Enzyme-Linked Immunosorbent Assay (ELISA). The plate has been pre-coated with Rat MCT4 antibody. MCT4 present in the sample is added and binds to antibodies coated on the wells. And then biotinylated Rat MCT4 Antibody is added and binds to MCT4 in the sample. Then Streptavidin-HRP is added and binds to the Biotinylated MCT4 antibody. After incubation unbound Streptavidin-HRP is washed away during a washing step. Substrate solution is then added and color develops in proportion to the amount of Rat MCT4. The reaction is terminated by addition of acidic stop solution and absorbance is measured at 450 nm.

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | [contact@aaabiotech.com](mailto:contact@aaabiotech.com)  
Online datasheet: <https://stage.aaabiotech.com/rat-elisa-kits/mast-cell-tryptase-4-mct4/122561>