

Human Complement Decay-Accelerating Factor, CD55 ELISA Kit | CD55 elisa kit

Human Complement Decay-Accelerating Factor, CD55 Primacu ELISA Kit

CATALOG #
AAA124292

REACTIVITY
Human


PRODUCT SYNONYMS

Complement Decay-Accelerating Factor; CD55; Human Complement Decay-Accelerating Factor; CD55 Primacu ELISA Kit; Complement decay-accelerating factor; CR; DAF; CD55 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Assay Type	Sandwich
Samples	Serum, plasma, cell culture supernates
Detection Range	23.44-1500pg/mL
Sensitivity	4.59pg/mL
Standard Solution	15ng/mL
Assay Time	3.5h
Preparation and Storage	Store at 2-8 degree C

IMAGES



Standard Curve (Sample)

Typical Testing Data/Standard Curve (for reference only)

ADDITIONAL INFORMATION

Related Product Information for CD55 elisa kit	CD55, also well known as decay-accelerating factor (DAF), is a member of the RCA (regulators of complement activation) family characterized by four to 30 SCRs (short consensus repeats) in their plasma-exposed regions. It is a major regulator of the alternative and classical pathways of complement activation and is expressed on all serum-exposed cells. CD55 is physiologically acting as an inhibitor of the complement system, but is also broadly expressed in malignant tumours. DAF seems to exert different functions beyond its immunological role such as promotion of tumorigenesis, decrease of complement mediated tumor cell lysis, autocrine loops for cell rescue and evasion of apoptosis, neoangiogenesis, invasiveness, cell motility. It is commonly hijacked by invading pathogens, including many enteroviruses and uropathogenic Escherichia coli, to promote cellular attachment prior to infection. This 70-75 kDa glycoprotein CD55 containing four SCR modules is involved in the regulation of the complement cascade. It inhibits complement activation by suppressing the function of C3/C5 convertases, thereby limiting local generation or deposition of C3a/C5a and membrane attack complex (MAC or C5b-9) production. DAF has been identified as a ligand for an activation-associated, seven-transmembrane lymphocyte receptor, CD97, which is a receptor mediating attachment and infection of several viruses and bacteria. In addition, it has been shown that DAF regulates the interplay between complement and T cell immunity in vivo, and thus may be implicated in immune and tumor biology.
Product Categories/Family for CD55 elisa kit	Primacu ELISA Kit; Immunology

NCBI AND UNIPROT INFORMATION

NCBI GI #	10835143
NCBI GeneID	1604
NCBI Accession #	NP_000565.1
NCBI GenBank Nucleotide #	NM_000574.3
UniProt Accession #	P08174; P09679; P78361; B1AP14; D3DT83; D3DT84
Molecular Weight	41,400 Da
NCBI Official Full Name	complement decay-accelerating factor isoform 1 preproprotein
NCBI Official Synonym Full Names	CD55 molecule, decay accelerating factor for complement (Cromer blood group)
NCBI Official Symbol	CD55
NCBI Official Synonym Symbols	CR; TC; DAF; CROM
NCBI Protein Information	complement decay-accelerating factor; CD55 antigen
UniProt Protein Name	Complement decay-accelerating factor
UniProt Gene Name	CD55
UniProt Synonym Gene Names	CR; DAF
UniProt Entry Name	DAF_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/complement-decay-accelerating-factor-cd55/124292>