

Human Serotransferrin, TF ELISA Kit | TF elisa kit

Human Serotransferrin, TF Primacu ELISA Kit

CATALOG #
AAA124332

REACTIVITY
Human


PRODUCT SYNONYMS

Serotransferrin; TF; Human Serotransferrin; TF Primacu ELISA Kit; Transferrin; TF elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Assay Type	Sandwich
Samples	Serum, plasma, cell culture supernates
Detection Range	15.63-1000pg/mL
Sensitivity	5.56pg/mL
Standard Solution	10ng/mL
Assay Time	3.5h
Preparation and Storage	Store at 2-8 degree C

IMAGES



Standard Curve (Sample)

Typical Testing Data/Standard Curve (for reference only)

ADDITIONAL INFORMATION

Related Product Information for TF elisa kit	Transferrin is a glycoprotein with an approximate molecular weight of 76.5 kDa. This glycoprotein is thought to have been created as a result of an ancient gene duplication event that led to generation of homologous C and N-terminal domains each of which binds one ion of ferric iron. The function of Transferrin is to transport iron from the intestine, reticuloendothelial system, and liver parenchymal cells to all proliferating cells in the body. This protein may also have a physiologic role as granulocyte/pollen-binding protein (GPBP) involved in the removal of certain organic matter and allergens from serum. Transferrins are iron binding transport proteins that bind Fe ³⁺ ion in association with the binding of an anion, usually bicarbonate. This transferrin binds only one Fe ³⁺ ion per protein molecule. Transports iron ions from the hemolymph into the eggs during the vitellogenic stage. Transferrins are iron binding transport proteins which can bind two Fe(3+) ions in association with the binding of an anion, usually bicarbonate. It is responsible for the transport of iron from sites of absorption and heme degradation to those of storage and utilization. Serum transferrin may also have a further role in stimulating cell proliferation. When a transferrin loaded with iron encounters with a transferring receptor on cell surface, transferring binds to it and, as a consequence, is transported into the cell in a vesicle by receptor-mediated endocytosis. The PH is reduced by hydrogen iron pumps. The lower pH causes transferrin to release its iron ions. The receptor is then transported through the endocytic cycle back to the cell surface, ready for another round of iron uptake. Each transferrin molecule has the ability to carry two iron ions in the ferric form.
Product Categories/Family for TF elisa kit	Primacu ELISA Kit; Cardiovascular; Signal transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	339453
NCBI GeneID	7018
UniProt Accession #	P02787; O43890; Q1HBA5; Q9NQB8; Q9UHV0
Molecular Weight	77,064 Da
NCBI Official Full Name	transferrin
NCBI Official Synonym Full Names	transferrin
NCBI Official Symbol	TF
NCBI Official Synonym Symbols	TFQTL1; PRO1557; PRO2086
NCBI Protein Information	serotransferrin; siderophilin; beta-1 metal-binding globulin
UniProt Protein Name	Serotransferrin
UniProt Gene Name	TF
UniProt Synonym Gene Names	Transferrin
UniProt Entry Name	TRFE_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/serotransferrin-tf/124332>