Rabbit Synaptophysin Monoclonal Antibody | anti-SYP antibody

Anti-Synaptophysin Rabbit Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA124474	**Rabbit**	**Human, Mouse, Rat**	**Western Blot, Immunocytochemistry, Immunohistochemistry, Immunofluorescence, Immunoprecipitation, Flow Cytometry, Functional Assay**


PRODUCT SYNONYMS

Synaptophysin; Monoclonal Antibody; Anti-Synaptophysin Rabbit Monoclonal Antibody; Major synaptic vesicle protein p38; SYP; anti-SYP antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	195
Applicable Applications for anti-SYP antibody	WB (Western Blot), ICC (Immunocytochemistry), IHC (Immunohistochemistry), IF (Immunofluorescence), IP (Immunoprecipitation), FCM/FACS (Flow Cytometry)
Immunogen	A synthesized peptide derived from human Synaptophysin
Subcellular Localization	Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane; Multi-pass membrane protein. Cell junction, synapse, synaptosome.
Tissue Specificity	Characteristic of a type of small (30-80 nm) neurosecretory vesicles, including presynaptic vesicles, but also vesicles of various neuroendocrine cells of both neuronal and epithelial phenotype.
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.

IMAGES



WB (Western Blot)
Western blot analysis of Synaptophysin expression in rat brain lysate (AAA124474).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-SYP monoclonal antibody overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for SYP



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded human kidney, using Synaptophysin Antibody (AAA124474)SYP was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-SYP Antibody (AAA124474)overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-SYP antibody	Possibly involved in structural functions as organizing other membrane components or in targeting the vesicles to the plasma membrane. Involved in the regulation of short-term and long-term synaptic plasticity (By similarity).

NCBI AND UNIPROT INFORMATION

NCBI GI #	27764867
NCBI GeneID	6855
NCBI Accession #	NP_003170.1
NCBI GenBank Nucleotide #	NM_003179.2
UniProt Accession #	P08247; Q6P2F7; B2R7L6; B7Z359
Molecular Weight	33845 MW
NCBI Official Full Name	synaptophysin
NCBI Official Synonym Full Names	synaptophysin
NCBI Official Symbol	SYP
NCBI Official Synonym Symbols	MRX96; MRXSYP
NCBI Protein Information	synaptophysin
UniProt Protein Name	Synaptophysin
UniProt Gene Name	**SYP**

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: https://stage.aaabiotech.com/monoclonal/synaptophysin-antibody-western-blot-immunocytochemistry-immunohistochemistry-immunofluorescence-immunoprecipitation-flow-cytometry-elisa/124474