

Rabbit Histone H2A Monoclonal Antibody | anti-HIST1H2AB antibody

Anti-Histone H2A Rabbit Monoclonal Antibody



CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA124511	Rabbit	Human, Mouse, Rat	Western Blot, Immunocytochemistry, Immunohistochemistry, Immunofluorescence

PRODUCT SYNONYMS

Histone H2A; Monoclonal Antibody; Anti-Histone H2A Rabbit Monoclonal Antibody; Histone H2A type 1-B/E; Histone H2A.2; Histone H2A/a; Histone H2A/m; HIST1H2AB; H2AFM; HIST1H2AE; H2AFA; anti-HIST1H2AB antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	130
Applicable Applications for anti-HIST1H2AB antibody	WB (Western Blot), ICC (Immunocytochemistry), IHC (Immunohistochemistry), IF (Immunofluorescence)
Immunogen	A synthesized peptide derived from human Histone H2A
Subcellular Localization	Nucleus. Chromosome.
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.



IHC (Immunohistochemistry)
Immunohistochemical analysis of paraffin-embedded human colon, using Histone H2A Antibody(AAA124511)HIST1H2AB was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-HIST1H2AB Antibody (AAA124511)overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

WB (Western Blot)
Western blot analysis of Histone H2A expression in HeLa cell lysate (AAA124511).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-HIST1H2AB monoclonal antibody overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for HIST1H2AB

ADDITIONAL INFORMATION

Related Product Information for anti-HIST1H2AB antibody	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Product Categories/Family for anti-HIST1H2AB antibody	Epigenetics and Nuclear Signaling; histones; h2a; crotonylated

NCBI AND UNIPROT INFORMATION

NCBI GI #	19557656
NCBI GeneID	8335
NCBI Accession #	NP_003504.2
NCBI GenBank Nucleotide #	NM_003513.2
UniProt Accession #	P04908; P28001; Q76P63
Molecular Weight	14135 MW
NCBI Official Full Name	histone H2A type 1-B/E
NCBI Official Synonym Full Names	histone cluster 1 H2A family member b
NCBI Official Symbol	HIST1H2AB
NCBI Official Synonym Symbols	H2A/m; H2AFM
NCBI Protein Information	histone H2A type 1-B/E
UniProt Protein Name	Histone H2A type 1-B/E
UniProt Gene Name	HIST1H2AB
UniProt Synonym Gene Names	H2AFM

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/histone-h2a-antibody-western-blot-immunocytochemistry-immunohistochemistry-immunofluorescence-elisa/124511>