

Rabbit Creb Monoclonal Antibody | anti-Creb antibody

Anti-Phospho-Creb (S133) Rabbit Monoclonal Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA124530	Rabbit	Human, Mouse, Rat	Western Blot, Immunocytochemistry, Immunohistochemistry, Immunofluorescence, Immunoprecipitation, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS

Creb; Monoclonal Antibody; Anti-Phospho-Creb (S133) Rabbit Monoclonal Antibody; Cyclic AMP-responsive element-binding protein 1; CREB-1; cAMP-responsive element-binding protein 1; CREB1; anti-Creb antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Purity/Purification	Affinity-chromatography
Form/Format	Liquid
Sequence Length	230
Applicable Applications for anti-Creb antibody	WB (Western Blot), ICC (Immunocytochemistry), IHC (Immunohistochemistry), IF (Immunofluorescence), IP (Immunoprecipitation), FCM/FACS (Flow Cytometry)
Immunogen	A synthesized peptide derived from human Phospho-Creb (S133)
Subcellular Localization	Nucleus.
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.



IHC (Immunohistochemistry)

Immunohistochemical analysis of paraffin-embedded human gastric carcinoma, using Phospho-Creb (S133) Antibody(AAA124530)CREB1 was detected in paraffin-embedded tissue section. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-CREB1 Antibody (AAA124530)overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

Related Product Information for anti-Creb antibody	Phosphorylation-dependent transcription factor that stimulates transcription upon binding to the DNA cAMP response element (CRE), a sequence present in many viral and cellular promoters. Transcription activation is enhanced by the TORC coactivators which act independently of Ser-133 phosphorylation. Involved in different cellular processes including the synchronization of circadian rhythmicity and the differentiation of adipose cells.
Product Categories/Family for anti-Creb antibody	Epigenetics and Nuclear Signaling ; Transcription ; Domain Families ; HLH/Leucine Zipper ; Epigenetics and Nuclear Signaling ; Transcription Factors ; Kits/Lysates/Other ; ELISA Kits ; signal transduction proteins ELISA Kits ; phosphoprotein ELISA Kits

NCBI GI #	4758054
NCBI GeneID	1385
NCBI Accession #	NP_004370.1
NCBI GenBank Nucleotide #	NM_004379.4
UniProt Accession #	P16220; P21934; Q6V963; Q9UMA7
Molecular Weight	36688 MW
NCBI Official Full Name	cyclic AMP-responsive element-binding protein 1 isoform A
NCBI Official Synonym Full Names	cAMP responsive element binding protein 1
NCBI Official Symbol	CREB1
NCBI Official Synonym Symbols	CREB; CREB-1
NCBI Protein Information	cyclic AMP-responsive element-binding protein 1
UniProt Protein Name	Cyclic AMP-responsive element-binding protein 1
UniProt Gene Name	CREB1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/creb-antibody-western-blot-immunocytochemistry-immunohistochemistry-immunofluorescence-immunoprecipitation-flow-cytometry-elisa/124530>