

Rabbit CCR4 Polyclonal Antibody | anti-CCR4 antibody

Anti-CCR4 Picoband Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA124592	Rabbit	Human, Mouse, Rat No cross reactivity with other proteins.	Western Blot

PRODUCT SYNONYMS

CCR4; Polyclonal Antibody; Anti-CCR4 Picoband Antibody; C-C chemokine receptor type 4; C-C CKR-4; CC-CKR-4; CCR-4; K5-5; CD194; CMKBR4; anti-CCR4 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat No cross reactivity with other proteins.
Clonality	Polyclonal
Purity/Purification	Immunogen affinity purified
Form/Format	Lyophilized
Sequence Length	360
Applicable Applications for anti-CCR4 antibody	WB (Western Blot)
Immunogen	A synthetic peptide corresponding to a sequence of human CCR4 (LVELEVLQDCTFERYLDAIQATETLAFVH).
Subcellular Localization	Cell membrane; Multi-pass membrane protein.
Tissue Specificity	Predominantly expressed in the thymus, in peripheral blood leukocytes, including T-cells, mostly CD4+ cells, and basophils, and in platelets; at lower levels, in the spleen and in monocytes. Detected also in macrophages, IL-2-activated natural killer cells and skin-homing memory T-cells, mostly the ones expressing the cutaneous lymphocyte antigen (CLA). Expressed in brain microvascular and coronary artery endothelial cells.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Contents	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.



WB (Western Blot)

Figure 1. Western blot analysis of CCR4 using anti-CCR4 antibody (AAA124592). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: rat thymus tissue lysates, Lane 2: mouse thymus tissue lysates, Lane 3: mouse NIH3T3 whole cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CCR4 antigen affinity purified polyclonal antibody at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for CCR4 at approximately 41KD. The expected band size for CCR4 is at 41KD.

ADDITIONAL INFORMATION

Related Product Information for anti-CCR4 antibody

CCR4, C-C chemokine receptor type 4 is a protein that in humans is encoded by the CCR4 gene. CCR4 has also recently been designated CD194 (cluster of differentiation 194). The protein encoded by this gene belongs to the G protein-coupled receptor family. It is a receptor for the following CC chemokines. The CCR4 gene is mapped to chromosome 3. Chemokines are a group of small structurally related proteins that regulate cell trafficking of various types of leukocytes. The chemokines also play fundamental roles in the development, homeostasis, and function of the immune system, and they have effects on cells of the central nervous system as well as on endothelial cells involved in angiogenesis or angiostasis.

Protein Function: High affinity receptor for the C-C type chemokines CCL17/TARC, CCL22/MDC and CKLF isoform 1/CKLF1. The activity of this receptor is mediated by G(i) proteins which activate a phosphatidylinositol-calcium second messenger system. Can function as a chemoattractant homing receptor on circulating memory lymphocytes and as a coreceptor for some primary HIV-2 isolates. In the CNS, could mediate hippocampal-neuron survival.

NCBI AND UNIPROT INFORMATION

NCBI GI #	5031627
NCBI GeneID	1233
NCBI Accession #	NP_005499.1
NCBI GenBank Nucleotide #	NM_005508.4
UniProt Accession #	P51679; Q9ULY6; Q9ULY7
NCBI Official Full Name	C-C chemokine receptor type 4
NCBI Official Synonym Full Names	C-C motif chemokine receptor 4
NCBI Official Symbol	CCR4
NCBI Official Synonym Symbols	CKR4; K5-5; CD194; CMKBR4; ChemR13; CC-CKR-4; HGCN:14099
NCBI Protein Information	C-C chemokine receptor type 4

UniProt Protein Name	C-C chemokine receptor type 4
UniProt Gene Name	CCR4
UniProt Synonym Gene Names	CMKBR4; C-C CKR-4; CC-CKR-4; CCR-4; CCR4

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ccr4-antibody-western-blot-elisa/124592>