

Rabbit AMPK alpha 1 Antibody | anti-AMPK alpha 1 antibody

Anti-AMPK alpha 1 antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA124608	Rabbit	Human, Mouse, Rat	ELISA, Western Blot

PRODUCT SYNONYMS

AMPK alpha 1; Anti-AMPK alpha 1 antibody; 5'-AMP-activated protein kinase catalytic subunit alpha-1; AMPK subunit alpha-1; anti-AMPK alpha 1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Isotype	IgG
Purity/Purification	Immunogen affinity purified
Form/Format	Lyophilized; Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Applicable Applications for anti-AMPK alpha 1 antibody	ELISA, WB (Western Blot)
Immunogen	E.Coli-derived human AMPK alpha 1 recombinant protein (Position: F377-R446).
Reconstitution	0.2ml of distilled water will yield a concentration of 500ug/ml
Preparation and Storage	Store at -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquoted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing



WB (Western Blot)

Figure 1. Western blot analysis of AMPK alpha 1 using anti-AMPK alpha 1 antibody (AAA124608). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human PANC-1 whole cell lysates, Lane 2: human Jurkat whole cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-AMPK alpha 1 antigen affinity purified polyclonal antibody at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for AMPK alpha 1 at approximately 63KD. The expected band size for AMPK alpha 1 is at 63KD.

ADDITIONAL INFORMATION

Related Product Information for anti-AMPK alpha 1 antibody	5'-AMP-activated protein kinase catalytic subunit alpha-1 is an enzyme that in humans is encoded by the PRKAA1 gene. The protein encoded by this gene belongs to the ser/thr protein kinase family. It is the catalytic subunit of the 5'-prime-AMP-activated protein kinase (AMPK). AMPK is a cellular energy sensor conserved in all eukaryotic cells. The kinase activity of AMPK is activated by the stimuli that increase the cellular AMP/ATP ratio. AMPK regulates the activities of a number of key metabolic enzymes through phosphorylation. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	94557301
NCBI GeneID	5562
NCBI Accession #	NP_006242.5
NCBI GenBank Nucleotide #	NM_006251.5
UniProt Accession #	Q13131; O00286; Q5D0E1; Q86VS1; Q9UNQ4; A8MTQ6; B2R7E1
NCBI Official Full Name	5'-AMP-activated protein kinase catalytic subunit alpha-1 isoform 1
NCBI Official Synonym Full Names	protein kinase AMP-activated catalytic subunit alpha 1
NCBI Official Symbol	PRKAA1
NCBI Official Synonym Symbols	AMPK; AMPKa1
NCBI Protein Information	5'-AMP-activated protein kinase catalytic subunit alpha-1
UniProt Protein Name	5'-AMP-activated protein kinase catalytic subunit alpha-1
UniProt Gene Name	PRKAA1
UniProt Synonym Gene Names	AMPK1; HMGCR kinase

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rabbit-human-mouse-rat-antibody/ampk-alpha-1/124608>