

Rabbit Factor D Polyclonal Antibody | anti-Cfd antibody

Anti-Factor D Picoband antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA124624	Rabbit	Human, Mouse, Rat No cross reactivity with other proteins.	Western Blot, Immunohistochemistry, ELISA

PRODUCT SYNONYMS


Factor D; Polyclonal Antibody; Anti-Factor D Picoband antibody; Complement factor D; Adipsin; C3 convertase activator; Endogenous vascular elastase; Properdin factor D; Cfd; Adn; Df; anti-Cfd antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat No cross reactivity with other proteins.
Clonality	Polyclonal
Form/Format	Lyophilized
Sequence Length	263
Applicable Applications for anti-Cfd antibody	WB (Western Blot), IHC (Immunohistochemistry), ELISA
Immunogen	E Coli-derived rat Factor D recombinant protein (Position: I26-A263).
Subcellular Localization	Secreted.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Contents	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.




IHC (Immunohistochemistry)
Figure 4. IHC analysis of Factor D using anti-Factor D antibody (AAA124624).Factor D was detected in paraffin-embedded section of rat small intestine tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-Factor D Antibody (AAA124624) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)
Figure 3. IHC analysis of Factor D using anti-Factor D antibody (AAA124624).Factor D was detected in paraffin-embedded section of rat lung tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-Factor D Antibody (AAA124624) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)
Figure 2. IHC analysis of Factor D using anti-Factor D antibody (AAA124624).Factor D was detected in paraffin-embedded section of mouse lung tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-Factor D Antibody (AAA124624) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)
Figure 1. Western blot analysis of Factor D using anti-Factor D antibody (AAA124624).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.Lane 1: rat RH35 cell lysates,Lane 2: rat PC-12 cell lysates.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Factor D antigen affinity purified polyclonal antibody at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for Factor D at approximately 27KD. The expected band size for Factor D is at 27KD.

ADDITIONAL INFORMATION

Related Product Information for anti-Cfd antibody	Description: Factor D is a protein which in humans is encoded by the CFD gene. The protein encoded by this gene is a member of the trypsin family of peptidases. The encoded protein is a component of the alternative complement pathway best known for its role in humoral suppression of infectious agents. It is also a serine protease that is secreted by adipocytes into the bloodstream. And it stimulates glucose transport for triglyceride accumulation in fats cells and inhibits lipolysis. Finally, the encoded protein has a high level of expression in fat, suggesting a role for adipose tissue in immune system biology. Protein Function: Factor D cleaves factor B when the latter is complexed with factor C3b, activating the C3bbb complex, which then becomes the C3 convertase of the alternate pathway. Its function is homologous to that of C1s in the classical pathway.
Product Categories/Family for anti-Cfd antibody	Stem Cells; Mesenchymal Stem Cells; chondrogenesis; Stem Cells; osteogenesis

NCBI AND UNIPROT INFORMATION

NCBI GI #	117647198
NCBI GeneID	54249
NCBI Accession #	NP_001071110.1
NCBI GenBank Nucleotide #	NM_001077642.1
UniProt Accession #	P32038
Molecular Weight	28,442 Da
NCBI Official Full Name	complement factor D
NCBI Official Synonym Full Names	complement factor D
NCBI Official Symbol	Cfd
NCBI Official Synonym Symbols	Df; Adn; EVE
NCBI Protein Information	complement factor D
UniProt Protein Name	Complement factor D
UniProt Gene Name	Cfd
UniProt Synonym Gene Names	Adn; Df

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/factor-d-antibody-western-blot-immunohistochemistry-elisa/124624>