

Rabbit KiSS1 receptor Polyclonal Antibody | anti-KISS1R antibody

Anti-KiSS1 receptor antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA124642	Rabbit	Human, Mouse, Rat No cross reactivity with other proteins.	Western Blot

PRODUCT SYNONYMS

KiSS1 receptor; Polyclonal Antibody; Anti-KiSS1 receptor antibody; KiSS-1 receptor; KiSS-1R; G-protein coupled receptor 54; G-protein coupled receptor OT7T175; hOT7T175; Hypogonadotropin-1; Kisspeptins receptor; Metastin receptor; KISS1R; AXOR12; GPR54; anti-KISS1R antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat No cross reactivity with other proteins.
Clonality	Polyclonal
Purity/Purification	Immunogen affinity purified
Form/Format	Lyophilized
Sequence Length	398
Applicable Applications for anti-KISS1R antibody	WB (Western Blot)
Immunogen	A synthetic peptide corresponding to a sequence of human KiSS1 receptor (MLRHLGRVAVRPAPADSALQGQVLAERAGAVRAK).
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Contents	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.



WB (Western Blot)

Figure 1. Western blot analysis of KiSS1 receptor using anti-KiSS1 receptor antibody (AAA124642). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human WISH whole cell lysates, Lane 2: human HepG2 whole cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-KiSS1 receptor antigen affinity purified polyclonal antibody at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for KiSS1 receptor at approximately 43KD. The expected band size for KiSS1 receptor is at 43KD.

NCBI AND UNIPROT INFORMATION

NCBI GI #	62865887
NCBI GeneID	84634
NCBI Accession #	NP_115940.2
NCBI GenBank Nucleotide #	NM_032551.4
UniProt Accession #	Q969F8; Q96QG0; A5D8U2; B2RTV1
NCBI Official Full Name	kiSS-1 receptor
NCBI Official Synonym Full Names	KISS1 receptor
NCBI Official Symbol	KISS1R
NCBI Official Synonym Symbols	HH8; CPPB1; GPR54; AXOR12; KISS-1R; HOT7T175
NCBI Protein Information	kiSS-1 receptor
UniProt Protein Name	KiSS-1 receptor
UniProt Gene Name	KISS1R
UniProt Synonym Gene Names	AXOR12; GPR54; KiSS-1R; hOT7T175

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.