

Rabbit CKB Polyclonal Antibody | anti-CKB antibody

Anti-CKB Picoband antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA124662	Rabbit	Human, Mouse, Rat No cross reactivity with other proteins.	Western Blot, Immunohistochemistry, ELISA

PRODUCT SYNONYMS


CKB; Polyclonal Antibody; Anti-CKB Picoband antibody; Creatine kinase B-type; B-CK; Creatine kinase B chain; CKBB; anti-CKB antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat No cross reactivity with other proteins.
Clonality	Polyclonal
Form/Format	Lyophilized
Sequence Length	381
Applicable Applications for anti-CKB antibody	WB (Western Blot), IHC (Immunohistochemistry), ELISA
Immunogen	E Coli-derived human CKB recombinant protein (Position: H7-K381).
Subcellular Localization	Cytoplasm.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Contents	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.



IHC (Immunohistochemistry)
Figure 4. IHC analysis of CKB using anti-CKB antibody (AAA124662).CKB was detected in paraffin-embedded section of rat kidney tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-CKB Antibody (AAA124662) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)
Figure 3. IHC analysis of CKB using anti-CKB antibody (AAA124662).CKB was detected in paraffin-embedded section of mouse kidney tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-CKB Antibody (AAA124662) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)
Figure 2. IHC analysis of CKB using anti-CKB antibody (AAA124662).CKB was detected in paraffin-embedded section of human colon cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-CKB Antibody (AAA124662) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)
Figure 1. Western blot analysis of CKB using anti-CKB antibody (AAA124662).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.Lane 1: human Hela whole cell lysates,Lane 2: human placenta tissue lysates,Lane 3: human COLO-320 whole cell lysates,Lane 4: human SW620 whole cell lysates,Lane 5: human MDA-MB-231 whole cell lysates,Lane 6: rat brain tissue lysates,Lane 7: mouse brain tissue lysates.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CKB antigen affinity purified polyclonal antibody at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for CKB at approximately 45KD. The expected band size for CKB is at 43KD.

ADDITIONAL INFORMATION

Related Product Information for anti-CKB antibody

Description: Brain-type creatine kinase also known as CK-BB is a creatine kinase that in humans is encoded by the CKB gene. The protein encoded by this gene is a cytoplasmic enzyme involved in energy homeostasis. The encoded protein reversibly catalyzes the transfer of phosphate between ATP and various phosphogens such as creatine phosphate. It acts as a homodimer in brain as well as in other tissues, and as a heterodimer with a similar muscle isozyme in heart. The encoded protein is a member of the ATP: guanido phosphotransferase protein family. A pseudogene of this gene has been characterized.

Protein Function: Reversibly catalyzes the transfer of phosphate between ATP and various phosphogens (e.g. creatine phosphate). Creatine kinase isoenzymes play a central role in energy transduction in tissues with large, fluctuating energy demands, such as skeletal muscle, heart, brain and spermatozoa.

NCBI AND UNIPROT INFORMATION

NCBI GI #	21536286
NCBI GeneID	1152
NCBI Accession #	NP_001814.2
NCBI GenBank Nucleotide #	NM_001823.4
UniProt Accession #	P12277; Q2LE07; Q6FG40; Q9UC66; A8K236; B2R5R4
Molecular Weight	42,644 Da
NCBI Official Full Name	creatine kinase B-type
NCBI Official Synonym Full Names	creatine kinase B
NCBI Official Symbol	CKB
NCBI Official Synonym Symbols	BCK; B-CK; CKBB; HEL-211; HEL-S-29
NCBI Protein Information	creatine kinase B-type
UniProt Protein Name	Creatine kinase B-type
UniProt Gene Name	CKB
UniProt Synonym Gene Names	CKBB; CPK-B

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ckb-antibody-western-blot-immunohistochemistry-elisa/124662>