

Rabbit FABP5 Polyclonal Antibody | anti-FABP5 antibody

Anti-FABP5 Picoband Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA124678	Rabbit	Human, Mouse, Rat No cross reactivity with other proteins.	Western Blot, Immunohistochemistry

PRODUCT SYNONYMS


FABP5; Polyclonal Antibody; Anti-FABP5 Picoband Antibody; Fatty acid-binding protein; epidermal; Epidermal-type fatty acid-binding protein; E-FABP; Fatty acid-binding protein 5; Psoriasis-associated fatty acid-binding protein homolog; PA-FABP; anti-FABP5 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat No cross reactivity with other proteins.
Clonality	Polyclonal
Form/Format	Lyophilized
Sequence Length	135
Applicable Applications for anti-FABP5 antibody	WB (Western Blot), IHC (Immunohistochemistry)
Immunogen	A synthetic peptide corresponding to a sequence of human FABP5 (KWRLMESHGFEEYMKELGVGLALRKMAAMAKPD).
Subcellular Localization	Cytoplasm.
Tissue Specificity	Keratinocytes; highly expressed in psoriatic skin.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Contents	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.




IHC (Immunohistochemistry)
Figure 5. IHC analysis of FABP5 using anti-FABP5 antibody (AAA124678).FABP5 was detected in paraffin-embedded section of rat lung tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-FABP5 Antibody (AAA124678) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC (Immunohistochemistry)
Figure 4. IHC analysis of FABP5 using anti-FABP5 antibody (AAA124678).FABP5 was detected in paraffin-embedded section of mouse lung tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-FABP5 Antibody (AAA124678) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)
Figure 3. IHC analysis of FABP5 using anti-FABP5 antibody (AAA124678).FABP5 was detected in paraffin-embedded section of human oesophagus squama cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-FABP5 Antibody (AAA124678) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)
Figure 2. IHC analysis of FABP5 using anti-FABP5 antibody (AAA124678).FABP5 was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-FABP5 Antibody (AAA124678) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of FABP5 using anti-FABP5 antibody (AAA124678). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human Hela whole cell lysates, Lane 2: human A549 whole cell lysates, Lane 3: rat thymus tissue lysates, Lane 4: mouse thymus tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-FABP5 antigen affinity purified polyclonal antibody at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for FABP5 at approximately 15KD. The expected band size for FABP5 is at 15KD.

ADDITIONAL INFORMATION

Related Product Information for anti-FABP5 antibody	Description: FABP5, Fatty acid-binding protein, epidermal, is a protein that in humans is encoded by the FABP5 gene. This gene encodes the fatty acid binding protein found in epidermal cells, and was first identified as being upregulated in psoriasis tissue. It is mapped to 8q21.13. Fatty acid binding proteins are a family of small, highly conserved, cytoplasmic proteins that bind long-chain fatty acids and other hydrophobic ligands. It is thought that FABPs roles include fatty acid uptake, transport, and metabolism. Protein Function: High specificity for fatty acids. Highest affinity for C18 chain length. Decreasing the chain length or introducing double bonds reduces the affinity. May be involved in keratinocyte differentiation.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557581
NCBI GeneID	2171
NCBI Accession #	NP_001435.1
NCBI GenBank Nucleotide #	NM_001444.2
UniProt Accession #	Q01469; B2R4K0
Molecular Weight	15,164 Da
NCBI Official Full Name	fatty acid-binding protein, epidermal
NCBI Official Synonym Full Names	fatty acid binding protein 5
NCBI Official Symbol	FABP5
NCBI Official Synonym Symbols	EFABP; KFABP; E-FABP; PAFABP; PA-FABP
NCBI Protein Information	fatty acid-binding protein, epidermal
UniProt Protein Name	Fatty acid-binding protein, epidermal
UniProt Gene Name	FABP5

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/fabp5-antibody-western-blot-immunohistochemistry-elisa/124678>