

Rabbit Frizzled 4 Polyclonal Antibody | anti-FZD4 antibody

Anti-Frizzled 4 Picoband antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA124680	Rabbit	Human, Mouse, Rat No cross reactivity with other proteins.	Western Blot, Immunohistochemistry

PRODUCT SYNONYMS

Frizzled 4; Polyclonal Antibody; Anti-Frizzled 4 Picoband antibody; Frizzled-4; Fz-4; hFz4; FzE4; CD344; FZD4; anti-FZD4 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat No cross reactivity with other proteins.
Clonality	Polyclonal
Form/Format	Lyophilized
Sequence Length	537
Applicable Applications for anti-FZD4 antibody	WB (Western Blot), IHC (Immunohistochemistry)
Immunogen	A synthetic peptide corresponding to a sequence of human Frizzled 4 (QNLGYNVTKMPNLVGHELQTDALQLTTFTPLIQY).
Subcellular Localization	Membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein.
Tissue Specificity	Almost ubiquitous. Largely expressed in adult heart, skeletal muscle, ovary, and fetal kidney. Moderate amounts in adult liver, kidney, pancreas, spleen, and fetal lung, and small amounts in placenta, adult lung, prostate, testis, colon, fetal brain and liver.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Contents	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Na ₃ .
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.




IHC (Immunohistochemistry)

Figure 5. IHC analysis of Frizzled 4 using anti-Frizzled 4 antibody (AAA124680).Frizzled 4 was detected in paraffin-embedded section of rat heart tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-Frizzled 4 Antibody (AAA124680) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 4. IHC analysis of Frizzled 4 using anti-Frizzled 4 antibody (AAA124680).Frizzled 4 was detected in paraffin-embedded section of mouse heart tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-Frizzled 4 Antibody (AAA124680) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 3. IHC analysis of Frizzled 4 using anti-Frizzled 4 antibody (AAA124680).Frizzled 4 was detected in paraffin-embedded section of human mammary cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-Frizzled 4 Antibody (AAA124680) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of Frizzled 4 using anti-Frizzled 4 antibody (AAA124680).Frizzled 4 was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-Frizzled 4 Antibody (AAA124680) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of Frizzled 4 using anti-Frizzled 4 antibody (AAA124680). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human placenta tissue lysates, Lane 2: human MCF-7 cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Frizzled 4 antigen affinity purified polyclonal antibody at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal was developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for Frizzled 4 at approximately 65KD. The expected band size for Frizzled 4 is at 60KD.

ADDITIONAL INFORMATION

Related Product Information for anti-FZD4 antibody	Description: Frizzled-4 is a protein that in humans is encoded by the FZD4 gene. This gene is a member of the frizzled gene family. Members of this family encode seven-transmembrane domain proteins that are receptors for the Wingless type MMTV integration site family of signaling proteins. Most frizzled receptors are coupled to the beta-catenin canonical signaling pathway. This protein may play a role as a positive regulator of the Wingless type MMTV integration site signaling pathway. A transcript variant retaining intronic sequence and encoding a shorter isoform has been described, however, its expression is not supported by other experimental evidence. Protein Function: Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin (CTNNB1) canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin (CTNNB1) and activation of Wnt target genes. Plays a critical role in retinal vascularization by acting as a receptor for Wnt proteins and norrin (NDP). In retina, it can be both activated by Wnt protein-binding, but also by a Wnt-independent signaling via binding of norrin (NDP), promoting in both cases beta-catenin (CTNNB1) accumulation and stimulation of LEF/TCF-mediated transcriptional programs. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues.
Product Categories/Family for anti-FZD4 antibody	Signal Transduction; Signaling Pathway; G Protein Signaling; gpcr; Epigenetics and Nuclear Signaling; Transcription; Domain Families; developmental families; Stem Cells; Signaling Pathways; wnt; Surface Molecules; Cancer; Oncoproteins/Suppressors; Tumor S

NCBI AND UNIPROT INFORMATION

NCBI GI #	22547161
NCBI GeneID	8322
NCBI Accession #	NP_036325.2
NCBI GenBank Nucleotide #	NM_012193.3
UniProt Accession #	Q9ULV1; Q14C97; Q6S9E4; A8K9Q3

Molecular Weight	59,881 Da
NCBI Official Full Name	frizzled-4
NCBI Official Synonym Full Names	frizzled class receptor 4
NCBI Official Symbol	FZD4
NCBI Official Synonym Symbols	Fz4; EVR1; FEVR; Fz-4; FzE4; GPCR; hFz4; CD344; FZD4S
NCBI Protein Information	frizzled-4
UniProt Protein Name	Frizzled-4
UniProt Gene Name	FZD4
UniProt Synonym Gene Names	Fz-4; hFz4

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/frizzled-4-antibody-western-blot-immunohistochemistry-elisa/124680>