

Rabbit MMP10 Polyclonal Antibody | anti-MMP10 antibody

Anti-MMP10 Picoband Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA124714	Rabbit	Human, Mouse, Rat No cross reactivity with other proteins.	Western Blot, Immunohistochemistry, ELISA

PRODUCT SYNONYMS

MMP10; Polyclonal Antibody; Anti-MMP10 Picoband Antibody; Stromelysin-2; SL-2; 3.4.24.22; Matrix metalloproteinase-10; MMP-10; Transin-2; STMY2; anti-MMP10 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat No cross reactivity with other proteins.
Clonality	Polyclonal
Purity/Purification	Immunogen affinity purified
Form/Format	Lyophilized
Concentration	Add 0.2ml of distilled water will yield a concentration of 500ug/ml. (varies by lot)
Sequence Length	476
Applicable Applications for anti-MMP10 antibody	WB (Western Blot), IHC (Immunohistochemistry), ELISA
Immunogen	E Coli-derived human MMP10 recombinant protein (Position: Y239-C476).
Subcellular Localization	Secreted, extracellular space, extracellular matrix.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.

IMAGES



IHC (Immunohistochemistry)

Figure 3. IHC analysis of MMP10 using anti-MMP10 antibody. MMP10 was detected in paraffin-embedded section of human mammary cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-MMP10 Antibody (A03759) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of MMP10 using anti-MMP10 antibody (A03759). MMP10 was detected in paraffin-embedded section of human colon cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-MMP10 Antibody (A03759) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of MMP10 using anti-MMP10 antibody (A03759). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: rat cardiac muscle tissue lysates, Lane 2: mouse cardiac muscle tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MMP10 antigen affinity purified polyclonal antibody at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for MMP10 at approximately 54KD. The expected band size for MMP10 is at 54KD.

ADDITIONAL INFORMATION

Related Product Information for anti-MMP10 antibody	Description: Stromelysin-2 also known as matrix metalloproteinase-10 (MMP-10) or transin-2 is an enzyme that in humans is encoded by the MMP10 gene. Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The enzyme encoded by this gene degrades proteoglycans and fibronectin. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3. Protein Function: Can degrade fibronectin, gelatins of type I, III, IV, and V; weakly collagens III, IV, and V. Activates procollagenase.
Product Categories/Family for anti-MMP10 antibody	Cardiovascular; Angiogenesis; Adhesion/ECM; Matrix Metalloproteinases; MMP; Cancer; Invasion/Microenvironment; ECM Enzymes; MMPS; Extracellular Matrix; Cell Biology; Proteolysis/Ubiquitin; Proteolytic Enzymes; Metalloprotease

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505205
NCBI GeneID	4319
NCBI Accession #	NP_002416.1
NCBI GenBank Nucleotide #	NM_002425.2
UniProt Accession #	P09238 ; Q53HH9 ; B2R9X9
Molecular Weight	54,151 Da
NCBI Official Full Name	stromelysin-2 preproprotein
NCBI Official Synonym Full Names	matrix metalloproteinase 10
NCBI Official Symbol	MMP10
NCBI Official Synonym Symbols	SL-2; STMY2
NCBI Protein Information	stromelysin-2
UniProt Protein Name	Stromelysin-2
UniProt Gene Name	MMP10
UniProt Synonym Gene Names	STMY2; SL-2; MMP-10

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/mmp10-antibody-western-blot-immunohistochemistry-elisa/124714>