

Rabbit TFF2 Polyclonal Antibody | anti-TFF2 antibody

Anti-TFF2 Picoband Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA124736	Rabbit	Mouse, Rat No cross reactivity with other proteins.	Western Blot, Immunohistochemistry, ELISA

PRODUCT SYNONYMS

TFF2; Polyclonal Antibody; Anti-TFF2 Picoband Antibody; Trefoil factor 2; Spasmolytic polypeptide; SP; Tff2; Sml1; anti-TFF2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Mouse, Rat No cross reactivity with other proteins.
Clonality	Polyclonal
Purity/Purification	Immunogen affinity purified
Form/Format	Lyophilized
Concentration	Add 0.2ml of distilled water will yield a concentration of 500ug/ml. (varies by lot)
Sequence Length	129
Applicable Applications for anti-TFF2 antibody	WB (Western Blot), IHC (Immunohistochemistry), ELISA
Immunogen	E Coli-derived rat TFF2 recombinant protein (Position: E24-Y129).
Subcellular Localization	Secreted.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of TFF2 using anti-TFF2 antibody (AAA124736).TFF2 was detected in paraffin-embedded section of rat gaster tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-TFF2 Antibody (AAA124736) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of TFF2 using anti-TFF2 antibody (AAA124736).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.Lane 1: rat gaster tissue lysates,Lane 2: mouse gaster tissue lysates.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TFF2 antigen affinity purified polyclonal antibody at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for TFF2 at approximately 14KD. The expected band size for TFF2 is at 14KD.

ADDITIONAL INFORMATION

Related Product Information for anti-TFF2 antibody	Description: Trefoil factor 2, also known as SP or SML1, is a protein that in humans is encoded by the TFF2 gene. This gene belongs to trefoil family, and members of the trefoil family may protect the mucosa from insults, stabilize the mucus layer, and affect healing of the epithelium. This gene is mapped to 21q22.3. TFF2 gene consists of 4 exons spanning 5.1 kb, with exons 2 and 3 encoding 1 trefoil domain each, and it is the only gene encoding 2 TFF-domains on separate exons. It has been found that the distribution of TFF2 and pS2 suggests involvement in repair-enhancing mechanisms. Protein Function: Inhibits gastrointestinal motility and gastric acid secretion. Could function as a structural component of gastric mucus, possibly by stabilizing glycoproteins in the mucus gel through interactions with carbohydrate side chains.
Product Categories/Family for anti-TFF2 antibody	Cell Biology; Other Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	16758702
NCBI GeneID	116592
NCBI Accession #	NP_446296.1
NCBI GenBank Nucleotide #	NM_053844.1
UniProt Accession #	Q09030
Molecular Weight	14,077 Da
NCBI Official Full Name	trefoil factor 2
NCBI Official Synonym Full Names	trefoil factor 2
NCBI Official Symbol	Tff2
NCBI Protein Information	trefoil factor 2

UniProt Protein Name	Trefoil factor 2
UniProt Gene Name	Tff2
UniProt Synonym Gene Names	Sml1; SP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/tff2-antibody-western-blot-immunohistochemistry-elisa/124736>