

Rabbit Cardiac FABP/FABP3 Polyclonal Antibody | anti-FABP3 antibody

Anti-Cardiac FABP/FABP3 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA124933	Rabbit	Human, Mouse, Rat	ELISA, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

Cardiac FABP/FABP3; Polyclonal Antibody; Anti-Cardiac FABP/FABP3 Antibody; Fatty acid-binding protein; heart; Fatty acid-binding protein 3; Heart-type fatty acid-binding protein; H-FABP; Mammary-derived growth inhibitor; MDGI; Muscle fatty acid-binding protein; M-FABP; FABP3; FABP11; Fatty acid binding protein 3; anti-FABP3 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	No cross reactivity with other proteins.
Purity/Purification	Immunogen Affinity Purified
Form/Format	Lyophilized
Sequence Length	133
Applicable Applications for anti-FABP3 antibody	ELISA (Direct ELISA), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived human Cardiac FABP recombinant protein (Position: V2-A133).
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Relevant Detection Systems	It is recommended to use an Enhanced Chemiluminescent Kit with anti-Rabbit IgG for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit for IHC-P.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.



IHC (Immunohistochemistry)

Figure 3. IHC analysis of Cardiac FABP using anti-Cardiac FABP antibody. Cardiac FABP was detected in paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/mL rabbit anti-Cardiac FABP antibody. Cardiac FABP was detected in paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/mL rabbit anti-Cardiac FABP antibody. Cardiac FABP was detected in paraffin-embedded section of mouse heart tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-Cardiac FABP antibody. Cardiac FABP was detected in paraffin-embedded section of rat heart tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-Cardiac FABP Antibody overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of Cardiac FABP using anti-Cardiac FABP antibody. Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: mouse heart tissue lysates, Lane 2: mouse heart tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Cardiac FABP antigen affinity purified polyclonal antibody at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for Cardiac FABP at approximately 15KD. The expected band size for Cardiac FABP is at 15KD.

ADDITIONAL INFORMATION

Related Product Information for anti-FABP3 antibody	Description: Rabbit IgG polyclonal antibody for Cardiac FABP detection. Tested with WB, IHC-P, Direct ELISA in Human; Mouse; Rat. Background: Heart-type fatty acid binding protein (hFABP), also known as mammary-derived growth inhibitor, is a protein that in humans is encoded by the FABP3 gene. The intracellular fatty acid-binding proteins (FABPs) belong to a multigene family. Fatty acid-binding protein 3 gene contains four exons and its function is to arrest growth of mammary epithelial cells. This gene is also a candidate tumor suppressor gene for human breast cancer. Cardiac-type fatty acid-binding protein (cFABP) from human heart muscle of three individuals was isolated and characterized as pl 5.3-cFABP.
References	1. "Entrez Gene: FABP3 fatty acid binding protein 3, muscle and heart (mammary-derived growthinhibitor)". Borchers, T., Hojrup, P., Nielsen, S. U., Roepstorff, P., Spener, F., Knudsen, J. Revision of the amino acid sequence of human heart fatty acid binding protein. Molec. Cell. Biochem. 98: 127-133, 1990. 3. Phelan CM, Larsson C, Baird S, Futreal PA, Ruttledge MH, Morgan K, Tonin P, Hung H, Korneluk RG, Pollak MN, Narod SA (Dec 1996). "The human mammary-derived growth inhibitor (MDGI) gene: genomic structure and mutation analysis in human breast tumors". Genomics 34 (1): 63-8.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758328
NCBI GeneID	2170
NCBI Accession #	NP_004093.1
NCBI GenBank Nucleotide #	NM_004102.4
UniProt Accession #	P05413; Q5VV93; Q99957; B2RAB6
Molecular Weight	14,858 Da
NCBI Official Full Name	fatty acid-binding protein, heart isoform 2
NCBI Official Synonym Full Names	fatty acid binding protein 3
NCBI Official Symbol	FABP3
NCBI Official Synonym Symbols	MDGI; FABP11; H-FABP; M-FABP; O-FABP
NCBI Protein Information	fatty acid-binding protein, heart

UniProt Protein Name	Fatty acid-binding protein, heart
UniProt Gene Name	FABP3
UniProt Synonym Gene Names	FABP11; MDGI; H-FABP; MDGI; M-FABP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cardiac-fabp-fabp3-antibody-direct-elisa-immunohistochemistry-western-blot-elisa/124933>