

Rabbit P cadherin/CDH3 Polyclonal Antibody | anti-CDH3 antibody

Anti-P cadherin/CDH3 Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA124952	Rabbit	Human, Mouse, Rat	ELISA, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

P cadherin/CDH3; Polyclonal Antibody; Anti-P cadherin/CDH3 Antibody; Cadherin-3; Placental cadherin; P-cadherin; CDH3; CDHP; Cadherin 3; type 1; P-cadherin (placental); anti-CDH3 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	No cross reactivity with other proteins.
Purity/Purification	Immunogen Affinity Purified
Form/Format	Lyophilized
Sequence Length	784
Applicable Applications for anti-CDH3 antibody	ELISA (Direct ELISA), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived human P cadherin recombinant protein (Position: Q126-H336).
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Relevant Detection Systems	It is recommended to use an Enhanced Chemiluminescent Kit with anti-Rabbit IgG for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit for IHC-P and IHC (F).
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.



IHC (Immunohistochemistry)

Figure 3. IHC analysis of P cadherin using anti-P cadherin antibody. P cadherin was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-P cadherin antibody. P cadherin was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-P cadherin antibody. P cadherin was detected in paraffin-embedded section of human placenta tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-P cadherin antibody. P cadherin was detected in paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-P cadherin antibody. P cadherin was detected in paraffin-embedded section of mouse intestine tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-P cadherin antibody overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of P cadherin using anti-P cadherin antibody. Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human Hela whole cell lysate, Lane 2: human K562 whole cell lysate, Lane 3: human A431 whole cell lysate, Lane 4: human Caco-2 whole cell lysate, Lane 5: rat brain tissue lysates, Lane 6: mouse brain tissue lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-P cadherin antigen affinity purified polyclonal antibody at 0.5 ug/ml overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for P cadherin at approximately 120KD. The expected band size for P cadherin is at 91KD.

ADDITIONAL INFORMATION

Related Product Information for anti-CDH3 antibody	Description: Rabbit IgG polyclonal antibody for P cadherin detection. Tested with WB, IHC-P, IHC-F, Direct ELISA in Human; Mouse; Rat. Background: Cadherins, such as CDH3, are integral membrane glycoproteins responsible for calcium-dependent cell-cell adhesion. Cadherin-3 is a protein that in humans is encoded by the CDH3 gene. This gene is a classical cadherin from the cadherin superfamily. The encoded protein is a calcium-dependent cell-cell adhesion glycoprotein composed of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. This gene is located in a six-cadherin cluster in a region on the long arm of chromosome 16 that is involved in loss of heterozygosity events in breast and prostate cancer. In addition, aberrant expression of this protein is observed in cervical adenocarcinomas. Mutations in this gene have been associated with congenital hypotrichosis with juvenile macular dystrophy.
References	1. Kaupmann K, Becker-Follmann J, Scherer G, Jockusch H, Starzinski-Powitz A (Dec 1992). "The gene for the cell adhesion molecule M-cadherin maps to mouse chromosome 8 and human chromosome 16q24.1-qter and is near the E-cadherin (uvomorulin) locus in both species". Genomics 14 (2): 488-90.

NCBI AND UNIPROT INFORMATION

NCBI GI #	953953841
NCBI GeneID	1001

NCBI Accession #	NP_001304124.1
NCBI GenBank Nucleotide #	NM_001317195.1
UniProt Accession #	P22223 ; Q05DI6 ; B2R6F4
Molecular Weight	86,552 Da
NCBI Official Full Name	cadherin-3 isoform 2
NCBI Official Synonym Full Names	cadherin 3
NCBI Official Symbol	CDH3
NCBI Official Synonym Symbols	CDHP; HJMD; PCAD
NCBI Protein Information	cadherin-3
UniProt Protein Name	Cadherin-3
UniProt Gene Name	CDH3
UniProt Synonym Gene Names	CDHP; P-cadherin

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/p-cadherin-cdh3-antibody-direct-elisa-immunohistochemistry-western-blot-elisa/124952>