

Human CD164 ELISA Kit | CD164 elisa kit

Human CD164 PicoKine ELISA Kit

CATALOG #
AAA125055

REACTIVITY
Human


PRODUCT SYNONYMS

CD164; Human CD164 PicoKine ELISA Kit; Human CD164; CD164 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	Natural and recombinant Human CD14
Sequence Length	197
Samples	Cell Culture Supernates, Serum And Plasma (Heparin, Edta, Citrate
Detection Range	156pg/ml-10000pg/ml
Sensitivity	<50pg/ml
Intra-assay Precision	Intra-Assay Precision (Precision within an assay): Three samples of known concentration were tested on one plate to assess intra-assay precision
Inter-assay Precision	Inter-Assay Precision (Precision accross assays):Three samples of known concentration were tested in separate assays to assess inter-assay precision.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CD164 elisa kit	Principle of the Assay: The Picokine Human CD14 Pre-Coated ELISA (Enzyme-Linked Immunosorbent Assay) kit is a solid phase immunoassay specially designed to measure Human CD14 with a 96-well strip plate that is pre-coated with antibody specific for CD14. The detection antibody is a biotinylated antibody specific for CD14. The capture antibody is A monoclonal antibody from mouse, the detection antibody is a biotinylated detection polyclonal antibody from goat. The kit contains recombinant Human CD14 with immunogen: Expression system for standard: CHO, Immunogen sequence: T20-C352. The kit is analytically validated with ready to use reagents. To measure Human CD14, add standards and samples to the wells, then add the biotinylated detection antibody. Wash the wells with PBS or TBS buffer, and add Avidin-Biotin-Peroxidase Complex (ABC-HRP). Wash away the unbounded ABC-HRP with PBS or TBS buffer and add TMB. TMB is substrate to HRP and will be catalyzed to produce a blue color product, which changes into yellow after adding acidic stop solution. The density of the yellow product is linearly propotional to Human CD14 in the sample. Read the density of the yellow product in each well using a plate reader, and benchmark the sample wells' readings against the standard curve to determine the concentration of Human CD14 in the sample!!Background/Introduction: CD14, Cluster of differentiation 14, ingle-copy gene encoding 2 protein forms: a 50-to 55-kD glycosylphosphatidylinositol-anchored membrane protein(mCD14) and a monocyte or liver-derived soluble serum protein(sCD14) that lacks the anchor. By in situ hybridization and study of somatic cell hybrid DNA that the gene is located at bands 5q23-q31. CD14 acts as a co-receptor(along with the Toll-like receptor TLR 4 and MD-2) for the detection of bacterial lipopolysaccharide(LPS). CD14 can bind LPS only in the presence of lipopolysaccharide-binding protein(LBP). Although LPS is considered its main ligand, CD14 also recognizes other pathogen-associated molecular patterns.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	9230741
NCBI GeneID	8763
NCBI Official Full Name	CD164
NCBI Official Synonym Full Names	CD164 molecule
NCBI Official Symbol	CD164
NCBI Official Synonym Symbols	DFNA66; MGC-24; MUC-24; MGC-24v; endolyn
NCBI Protein Information	sialomucin core protein 24

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/cd164/125055>