

Mouse anti-Human Cytokeratin 19 Monoclonal Antibody | anti-KRT19 antibody

Anti-Cytokeratin 19 Antibody (monoclonal, 3D4)



CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA125147	Mouse	Human	Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

Cytokeratin 19; Monoclonal Antibody; Anti-Cytokeratin 19 Antibody (monoclonal; 3D4); Keratin; type I cytoskeletal 19; Cytokeratin-19; CK-19; Keratin-19; K19; KRT19; Keratin 19; anti-KRT19 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Clone Number	3D4
Specificity	No cross reactivity with other proteins.
Purity/Purification	Immunogen Affinity Purified
Form/Format	Lyophilized
Sequence Length	400
Applicable Applications for anti-KRT19 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Cytokeratin 19 (334-372aa QLAHIQALISGIEAQLGDVRADSERQNQEYQRLMDIKSR), different from the related mouse and rat sequences by nine amino acids.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Relevant Detection Systems	It is recommended to use an Enhanced Chemiluminescent Kit with anti-Rabbit IgG for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit for IHC-P.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.




IF (Immunofluorescence)

Figure 3. IF analysis of Cytokeratin 19 using anti- Cytokeratin 19 antibody (M02101-2).Cytokeratin 19 was detected in paraffin-embedded section of human intestinal cancer tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml mouse anti- Cytokeratin 19 Antibody (M02101-2) overnight at 4 degree C. DyLight550 Conjugated Goat Anti-Mouse IgG was used as secondary antibody at 1:200 dilution and incubated for 30 minutes at 37 degree C. Visualize using a fluorescence microscope and filter sets appropriate for the label used.

IF (Immunofluorescence)

Figure 3. IF analysis of Cytokeratin 19 using anti- Cytokeratin 19 antibody (M02101-2).Cytokeratin 19 was detected in paraffin-embedded section of human intestinal cancer tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml mouse anti- Cytokeratin 19 Antibody (M02101-2) overnight at 4 degree C. DyLight550 Conjugated Goat Anti-Mouse IgG was used as secondary antibody at 1:200 dilution and incubated for 30 minutes at 37 degree C. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



WB (Western Blot)

Figure 1. Western blot analysis of Cytokeratin 19 using anti-Cytokeratin 19 antibody (M02101-2).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.Lane 1: human HeLa whole cell lysates,Lane 2: human placenta tissue lysates,Lane 3: human SK-OV-3 whole cell lysates,Lane 4: human COLO-320 whole cell lysates.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-Cytokeratin 19 antigen affinity purified monoclonal antibody at 0.5 ug/ml overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a Biotin Conjugated goat anti-mouse IgG secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system.

ADDITIONAL INFORMATION

Related Product Information for anti-KRT19 antibody

Description: Mouse IgG monoclonal antibody for Cytokeratin 19 detection. Tested with WB, IHC-P, IF in Human. Background: Keratin, type I cytoskeletal 19 is a protein that in humans is encoded by the KRT19 gene. The protein encoded by this gene is a member of the keratin family. It is specifically expressed in the periderm, the transiently superficial layer that envelops the developing epidermis. The type I cytokeratins are clustered in a region of chromosome 17q12-q21. Due to its high sensitivity, KRT19 is the most used marker for the RT-PCR-mediated detection of tumor cells disseminated in lymph nodes, peripheral blood, and bone marrow of breast cancer patients. Keratin 19 is often used together with keratin 8 and keratin 18 to differentiate cells of epithelial origin from hematopoietic cells in tests that enumerate circulating tumor cells in blood.

References	1. "Entrez Gene: KRT19 keratin 19". 2. Schweizer J, Bowden PE, Coulombe PA, Langbein L, Lane EB, Magin TM, Maltais L, Omary MB, Parry DA, Rogers MA, Wright MW (Jul 2006). "New consensus nomenclature for mammalian keratins". J Cell Biol 174 (2): 169-74. 3. W. Jeffrey Allard, Jeri Matera, M. Craig Miller, et al. (October 2004). "Tumor Cells Circulate in the Peripheral Blood of All Major Carcinomas but not in Healthy Subjects or Patients With Nonmalignant Diseases". Clin. Cancer Research 10 (20): 6897-6904.
------------	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	24234699
NCBI GeneID	3880
NCBI Accession #	NP_002267.2
NCBI GenBank Nucleotide #	NM_002276.5
UniProt Accession #	P08727; Q5XG83; Q6NW33; Q7L5M9; Q96A53; Q96FV1; Q9BYF9; Q9P1Y4; B2R874
Molecular Weight	44,106 Da
NCBI Official Full Name	keratin, type I cytoskeletal 19
NCBI Official Synonym Full Names	keratin 19
NCBI Official Symbol	KRT19
NCBI Official Synonym Symbols	K19; CK19; K1CS
NCBI Protein Information	keratin, type I cytoskeletal 19
UniProt Protein Name	Keratin, type I cytoskeletal 19
UniProt Gene Name	KRT19
UniProt Synonym Gene Names	CK-19; K19

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/monoclonal/cytokeratin-19-antibody-clone-3d4-immunofluorescence-immunohistochemistry-western-blot-elisa/125147>