

Rabbit anti-Human P Glycoprotein/ABCB1 Polyclonal Antibody | anti-ABCB1 antibody

Anti-P Glycoprotein/ABCB1 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA125948	Rabbit	Human	Flow Cytometry, Functional Assay, Immunofluorescence, Immunocytochemistry, Western Blot

PRODUCT SYNONYMS

P Glycoprotein/ABCB1; Polyclonal Antibody; Anti-P Glycoprotein/ABCB1 Antibody Picoband; anti-ABCB1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-ABCB1 antibody	FCM/FACS (Flow Cytometry), IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	A synthetic peptide corresponding to a sequence of human P Glycoprotein/ABCB1 (TKEALDESIPPVSFWRMK).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




FCM/FACS (Flow Cytometry)

Figure 3. Flow Cytometry analysis of 293T cells using anti-P Glycoprotein/ABCB1 antibody (AAA125948).Overlay histogram showing 293T cells stained with AAA125948 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-P Glycoprotein/ABCB1 Antibody (AAA125948, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IF (Immunofluorescence)

Figure 2. IF analysis of P Glycoprotein/ABCB1 using anti-P Glycoprotein/ABCB1 antibody (AAA125948).P Glycoprotein/ABCB1 was detected in an immunocytochemical section of MCF-7 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-P Glycoprotein/ABCB1 Antibody (AAA125948) overnight at 4 degree C. DyLight488 Conjugated Goat Anti-Rabbit IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



WB (Western Blot)

Figure 1. Western blot analysis of P Glycoprotein/ABCB1 using anti-P Glycoprotein/ABCB1 antibody (AAA125948).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human HepG2 whole cell lysates,Lane 2: human Caco-2 whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-P Glycoprotein/ABCB1 antigen affinity purified polyclonal antibody (#AAA125948) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for P Glycoprotein/ABCB1 at approximately 141 kDa. The expected band size for P Glycoprotein/ABCB1 is at 141 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-ABCB1 antibody

P-GP, also called ABCB1 or PGY1, is a glycoprotein that in humans is encoded by the ABCB1 gene. It is mapped to 7q21.12. P-GP is a well-characterized ABC-transporter (which transports a wide variety of substrates across extra- and intracellular membranes) of the MDR/TAP subfamily. It is an important protein of the cell membrane that pumps many foreign substances out of cells. More formally, it is an ATP-dependent drug efflux pump with broad substrate specificity. P-GP is an ATP-dependent drug efflux pump for xenobiotic compounds with broad substrate specificity. It is responsible for decreased drug accumulation in multidrug-resistant cells and often mediates the development of resistance to anticancer drugs. This protein also functions as a transporter in the blood-brain barrier.

Product Categories/Family for anti-
ABCB1 antibody

Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	42741659
NCBI GeneID	5243
NCBI Accession #	NP_000918.2
NCBI GenBank Nucleotide #	NM_000927.4
UniProt Accession #	P08183; Q12755; Q14812; A8K294
Molecular Weight	141kDa
NCBI Official Full Name	multidrug resistance protein 1
NCBI Official Synonym Full Names	ATP-binding cassette, sub-family B (MDR/TAP), member 1
NCBI Official Symbol	ABCB1
NCBI Official Synonym Symbols	CLCS; MDR1; P-GP; PGY1; ABC20; CD243; GP170
NCBI Protein Information	multidrug resistance protein 1; P-glycoprotein 1; colchicin sensitivity; doxorubicin resistance
UniProt Protein Name	Multidrug resistance protein 1
UniProt Gene Name	ABCB1
UniProt Synonym Gene Names	MDR1; PGY1
UniProt Entry Name	MDR1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/p-glycoprotein-abcb1-antibody-flow-cytometry-immunofluorescence-immunocytochemistry-western-blot-elisa/125948>