

Rabbit anti-Mouse, Rat IL13 Polyclonal Antibody | anti-IL13 antibody

Anti-IL13 Antibody Picoband

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA125951	Rabbit	Mouse, Rat	ELISA, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

IL13; Polyclonal Antibody; Anti-IL13 Antibody Picoband; anti-IL13 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-IL13 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived mouse IL13 recombinant protein (Position: P22-F131).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.

IMAGES




IHC (Immunohistochemistry)

Figure 3. IHC analysis of IL13 using anti-IL13 antibody (AAA125951). IL13 was detected in a paraffin-embedded section of rat lymph node tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-IL13 Antibody (AAA125951) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of IL13 using anti-IL13 antibody (AAA125951). IL13 was detected in a paraffin-embedded section of mouse spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-IL13 Antibody (AAA125951) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of IL13 using anti-IL13 antibody (AAA125951). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: mouse spleen tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-IL13 antigen affinity purified polyclonal antibody (#AAA125951) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for IL13 at approximately 19 kDa. The expected band size for IL13 is at 19 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-IL13 antibody	Interleukin 13 is a protein that in humans is encoded by the IL-13 gene. It is a kind of cytokine secreted by many cell types, but especially T helper type 2 (Th2) cells, which is an important mediator of allergic inflammation and disease. The IL-13 gene is mapped to 5q23-q31. IL-13 induces its effects through a multi-subunit receptor that includes the alpha chain of the IL-4 receptor (IL-4Ralpha) and at least one of two known IL-13-specific binding chains. Furthermore, this gene acts more prominently as a molecular bridge linking allergic inflammatory cells to the non-immune cells in contact with them, thereby altering physiological function.
Product Categories/Family for anti-IL13 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	6680403
NCBI GeneID	16163
NCBI Accession #	NP_032381.1
NCBI GenBank Nucleotide #	NM_008355.3
UniProt Accession #	P20109
Molecular Weight	14,108 Da
NCBI Official Full Name	interleukin-13
NCBI Official Synonym Full Names	interleukin 13
NCBI Official Symbol	IL13
NCBI Official Synonym Symbols	IL-13
NCBI Protein Information	interleukin-13; T-cell activation protein P600
UniProt Protein Name	Interleukin-13
UniProt Gene Name	IL13

UniProt Synonym Gene Names	IL-13; IL-13
UniProt Entry Name	IL13_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/il13-antibody-immunohistochemistry-western-blot-elisa/125951>