

Rabbit anti-Human ICAM1 Polyclonal Antibody | anti-ICAM1 antibody

Anti-ICAM1 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA125971	Rabbit	Human	ELISA, Flow Cytometry, Functional Assay, Western Blot

PRODUCT SYNONYMS


ICAM1; Polyclonal Antibody; Anti-ICAM1 Antibody Picoband; anti-ICAM1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-ICAM1 antibody	ELISA, FCM/FACS (Flow Cytometry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human ICAM1 recombinant protein (Position: D53-K519).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)
Figure 2. Flow Cytometry analysis of HepG2 cells using anti-ICAM1 antibody (AAA125971).Overlay histogram showing HepG2 cells stained with AAA125971 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-ICAM1 Antibody (AAA125971, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)
Figure 1. Western blot analysis of ICAM1 using anti-ICAM1 antibody (AAA125971).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human Ramos whole cell lysates,Lane 2: human Raji whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ICAM1 antigen affinity purified polyclonal antibody (#AAA125971) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for ICAM1 at approximately 95 kDa. The expected band size for ICAM1 is at 95 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-ICAM1 antibody	CD54, also known as ICAM-1. Intercellular adhesion molecule-1 (ICAM1) is a ligand for lymphocyte function-associated (LFA) antigens. ICAM-1 is an integral membrane protein, a member of the immunoglobulin superfamily, and a ligand for LFA-1, a beta 2 leukocyte integrin. This protein is the major human rhinovirus receptor. The ICAM1 gene is mapped to human chromosome 19. In humans, lymphocyte adhesion to cells is mediated by the protein heterodimer CD11a/CD18 (Leu-CAMa, LFA-1) and its ligand CD54 (ICAM-1).
Product Categories/Family for anti-ICAM1 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	167466198
NCBI GeneID	3383
NCBI Accession #	NP_000192.2
NCBI GenBank Nucleotide #	NM_000201.2
UniProt Accession #	P05362; Q5NKV7; Q96B50; B2R6M3
Molecular Weight	57,825 Da
NCBI Official Full Name	intercellular adhesion molecule 1
NCBI Official Synonym Full Names	intercellular adhesion molecule 1
NCBI Official Symbol	ICAM1
NCBI Official Synonym Symbols	BB2; CD54; P3.58
NCBI Protein Information	intercellular adhesion molecule 1; ICAM-1; cell surface glycoprotein P3.58; major group rhinovirus receptor; intercellular adhesion molecule 1 (CD54), human rhinovirus receptor

UniProt Protein Name	Intercellular adhesion molecule 1
UniProt Gene Name	ICAM1
UniProt Synonym Gene Names	ICAM-1
UniProt Entry Name	ICAM1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/icam1-antibody-flow-cytometry-western-blot-elisa/125971>