

Rabbit anti-Human, Mouse Cd80 Polyclonal Antibody | anti-CD80 antibody

Anti-Cd80 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA125979	Rabbit	Human, Mouse	Western Blot, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS


Cd80; Polyclonal Antibody; Anti-Cd80 Antibody Picoband; anti-CD80 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-CD80 antibody	WB (Western Blot), FCM/FACS (Flow Cytometry), ELISA
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived mouse Cd80 recombinant protein (Position: D62-K92).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)
Figure 2. Flow Cytometry analysis of ANA-1 cells using anti-Cd80 antibody (AAA125979).Overlay histogram showing ANA-1 cells stained with AAA125979 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Cd80 Antibody (AAA125979, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)
Figure 1. Western blot analysis of Cd80 using anti-Cd80 antibody (AAA125979).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human Daudi whole cell lysates,Lane 2: human K562 whole cell lysates,Lane 3: mouse RAW264.7 whole cell lysates,Lane 4: mouse ANA-1 whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Cd80 antigen affinity purified polyclonal antibody (#AAA125979) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for Cd80 at approximately 70 kDa. The expected band size for Cd80 is at 33 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-CD80 antibody	Cluster of Differentiation 80 (also CD80 and B7-1) is a protein found on activated B cells and monocytes that provides a costimulatory signal necessary for T cell activation and survival. It is the ligand for two different proteins on the T cell surface: CD28 (for autoregulation and intercellular association) and CTLA-4 (for attenuation of regulation and cellular disassociation). CD80 works in tandem with CD86 to prime T cells. The CD80 genes encode B7-1 which are structurally similar members of the immunoglobulin superfamily expressed on a variety of hematopoietic cell types.
Product Categories/Family for anti-CD80 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	112983684
NCBI GeneID	12519
NCBI Accession #	NP_033985.3
NCBI GenBank Nucleotide #	NM_009855.2
UniProt Accession #	Q00609; Q61332
Molecular Weight	24,051 Da
NCBI Official Full Name	T-lymphocyte activation antigen CD80
NCBI Official Synonym Full Names	CD80 antigen
NCBI Official Symbol	Cd80
NCBI Official Synonym Symbols	B71; Ly53; TSA1; Cd28l; Ly-53; MIC17
NCBI Protein Information	T-lymphocyte activation antigen CD80

UniProt Protein Name	T-lymphocyte activation antigen CD80
UniProt Gene Name	Cd80
UniProt Synonym Gene Names	B7; B7

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd80-antibody-western-blot-flow-cytometry-elisa/125979>