

Rabbit anti-Human Ki67/MKI67 Polyclonal Antibody | anti-MKI67 antibody

Anti-Ki67/MKI67 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA125994	Rabbit	Human	ELISA, Immunohistochemistry

PRODUCT SYNONYMS


Ki67/MKI67; Polyclonal Antibody; Anti-Ki67/MKI67 Antibody Picoband; anti-MKI67 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-MKI67 antibody	ELISA, IHC (Immunohistochemistry)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human Ki67/MKI67 recombinant protein (Position: R5-Q2358).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




IHC (Immunohistochemistry)
Figure 5. IHC analysis of Ki67/MKI67 using anti-Ki67/MKI67 antibody (AAA125994).Ki67/MKI67 was detected in a paraffin-embedded section of human laryngeal squamous cell carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Ki67/MKI67 Antibody (AAA125994) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.




IHC (Immunohistochemistry)
Figure 4. IHC analysis of Ki67/MKI67 using anti-Ki67/MKI67 antibody (AAA125994).Ki67/MKI67 was detected in a paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Ki67/MKI67 Antibody (AAA125994) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)
Figure 3. IHC analysis of Ki67/MKI67 using anti-Ki67/MKI67 antibody (AAA125994).Ki67/MKI67 was detected in a paraffin-embedded section of human renal carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Ki67/MKI67 Antibody (AAA125994) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)
Figure 2. IHC analysis of Ki67/MKI67 using anti-Ki67/MKI67 antibody (AAA125994).Ki67/MKI67 was detected in a paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Ki67/MKI67 Antibody (AAA125994) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 1. IHC analysis of Ki67/MKI67 using anti-Ki67/MKI67 antibody (AAA125994). Ki67/MKI67 was detected in a paraffin-embedded section of human colonic adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-Ki67/MKI67 Antibody (AAA125994) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-MKI67 antibody	Ki-67(Proliferation-related Ki-67 antigen), also known as MKI67 or KIA, is a protein that in humans is encoded by the MKI67 gene. From study of a panel of human-rodent somatic cell hybrids, it has been demonstrated that a gene involved in the expression of the MKI67 antigen is located on chromosome 10. By in situ hybridization, Fonatsch et al. (1991) regionalized the MKI67 gene to chromosome 10q25-qter. By FISH, Traut et al. (1998) mapped the mouse Mki67 gene to chromosome 7F3-F5. Antigen KI-67 is a nuclear protein that is associated with and may be necessary for cellular proliferation. Furthermore it is associated with ribosomal RNA transcription. Inactivation of antigen KI-67 leads to inhibition of ribosomal RNA synthesis.
Product Categories/Family for anti-MKI67 antibody	Primary Antibodies , Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	225543215
NCBI GeneID	4288
NCBI Accession #	NP_001139438.1
NCBI GenBank Nucleotide #	NM_001145966.1
UniProt Accession #	P46013 ; Q5VWH2
Molecular Weight	358,694 Da
NCBI Official Full Name	antigen KI-67 isoform 2
NCBI Official Synonym Full Names	antigen identified by monoclonal antibody Ki-67
NCBI Official Symbol	MKI67
NCBI Official Synonym Symbols	KIA
NCBI Protein Information	antigen KI-67; proliferation-related Ki-67 antigen
UniProt Protein Name	Antigen KI-67
UniProt Gene Name	MKI67
UniProt Entry Name	KI67_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/ki67-mki67-antibody-immunohistochemistry-elisa/125994>