

Rabbit NAK/TBK1 Polyclonal Antibody | anti-TBK1 antibody

Anti-NAK/TBK1 Antibody Picoband

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA125997	Rabbit	Human, Mouse, Rat	ELISA, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

NAK/TBK1; Polyclonal Antibody; Anti-NAK/TBK1 Antibody Picoband; anti-TBK1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-TBK1 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human NAK/TBK1 recombinant protein (Position: M1-L729).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.

IMAGES



IHC (Immunohistochemistry)

Figure 2. IHC analysis of NAK/TBK1 using anti-NAK/TBK1 antibody (AAA125997).NAK/TBK1 was detected in a paraffin-embedded section of mouse testis tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-NAK/TBK1 Antibody (AAA125997) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of NAK/TBK1 using anti-NAK/TBK1 antibody (AAA125997).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human 293T whole cell lysates,Lane 2: human HepG2 whole cell lysates,Lane 3: human LNCAP whole cell lysates,Lane 4: rat brain tissue lysates,Lane 5: rat C6 whole cell lysates,Lane 7: mouse brain tissue lysates,Lane 8: mouse Neuro-2a whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-NAK/TBK1 antigen affinity purified polyclonal antibody (#AAA125997) at 0.25 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for NAK/TBK1 at approximately 84 kDa. The expected band size for NAK/TBK1 is at 84 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-TBK1 antibody	Serine/threonine-protein kinase TBK1, also called TANK-binding kinase 1 or NF-kappa-B-activating kinase is an enzyme that in humans is encoded by the TBK1 gene. The gene was assigned to human chromosome 12q14.2. Serine/threonine kinase plays an essential role in regulating inflammatory responses to foreign agents. TBK1 and NF-kappa-B signaling are essential in KRAS mutant tumors, and established a general approach for the rational identification of codependent pathways in cancer.
Product Categories/Family for anti-TBK1 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	7019547
NCBI GeneID	29110
NCBI Accession #	NP_037386.1
NCBI GenBank Nucleotide #	NM_013254.3
UniProt Accession #	Q9UHD2; Q8IYV3; Q9NUJ5; A8K4S4
Molecular Weight	84kDa
NCBI Official Full Name	serine/threonine-protein kinase TBK1
NCBI Official Synonym Full Names	TANK-binding kinase 1
NCBI Official Symbol	TBK1
NCBI Official Synonym Symbols	NAK; T2K
NCBI Protein Information	serine/threonine-protein kinase TBK1; NF-kB-activating kinase; NF-kappa-B-activating kinase
UniProt Protein Name	Serine/threonine-protein kinase TBK1
UniProt Gene Name	TBK1
UniProt Synonym Gene Names	NAK
UniProt Entry Name	TBK1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/nak-tbk1-antibody-immunohistochemistry-western-blot-elisa/125997>