

Rabbit anti-Human ATR Polyclonal Antibody | anti-ATR antibody

Anti-ATR Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA125998	Rabbit	Human	ELISA, Flow Cytometry, Functional Assay, Western Blot

PRODUCT SYNONYMS


ATR; Polyclonal Antibody; Anti-ATR Antibody Picoband; anti-ATR antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-ATR antibody	ELISA, FCM/FACS (Flow Cytometry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human ATR recombinant protein (Position: E730-A1303).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)
Figure 2. Flow Cytometry analysis of U87 cells using anti-ATR antibody (AAA125998). Overlay histogram showing U87 cells stained with AAA125998 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-ATR Antibody (AAA125998, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)
Figure 1. Western blot analysis of ATR using anti-ATR antibody (AAA125998). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HeLa whole cell lysates, Lane 2: human Daudi whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ATR antigen affinity purified polyclonal antibody (#AAA125998) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for ATR at approximately 301 kDa. The expected band size for ATR is at 301 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-ATR antibody	Serine/threonine-protein kinase ATR also known as ataxia telangiectasia and Rad3-related protein (ATR) or FRAP-related protein 1 (FRP1) is an enzyme that, in humans, is encoded by the ATR gene. The protein encoded by this gene is a serine/threonine kinase and DNA damage sensor, activating cell cycle checkpoint signaling upon DNA stress. The encoded protein can phosphorylate and activate several proteins involved in the inhibition of DNA replication and mitosis, and can promote DNA repair, recombination, and apoptosis. This protein is also important for fragile site stability and centrosome duplication. Defects in this gene are a cause of Seckel syndrome 1.
Product Categories/Family for anti-ATR antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	157266317
NCBI GeneID	545
NCBI Accession #	NP_001175.2
NCBI GenBank Nucleotide #	NM_001184.3
UniProt Accession #	Q13535; Q59HB2; Q7KYL3; Q93051; Q9BXK4
Molecular Weight	301,367 Da
NCBI Official Full Name	serine/threonine-protein kinase ATR
NCBI Official Synonym Full Names	ataxia telangiectasia and Rad3 related
NCBI Official Symbol	ATR
NCBI Official Synonym Symbols	FRP1; MEC1; SCKL; FCTCS; SCKL1
NCBI Protein Information	serine/threonine-protein kinase ATR; Rad3 related protein; FRAP-related protein 1; FRAP-related protein-1; MEC1, mitosis entry checkpoint 1, homolog; ataxia telangiectasia and Rad3-related protein

UniProt Protein Name	Serine/threonine-protein kinase ATR
UniProt Gene Name	ATR
UniProt Synonym Gene Names	FRP1
UniProt Entry Name	ATR_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/atr-antibody-flow-cytometry-western-blot-elisa/125998>