

Rabbit anti-Human, Rat MEK1/MAP2K1 Polyclonal Antibody | anti-MAP2K1 antibody

Anti-MEK1/MAP2K1 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126001	Rabbit	Human, Rat	Western Blot, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS


MEK1/MAP2K1; Polyclonal Antibody; Anti-MEK1/MAP2K1 Antibody Picoband; anti-MAP2K1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-MAP2K1 antibody	WB (Western Blot), FCM/FACS (Flow Cytometry), ELISA
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human MEK1/MAP2K1 recombinant protein (Position: M1-V393).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)
Figure 2. Flow Cytometry analysis of 293T cells using anti-MEK1/MAP2K1 antibody (AAA126001).Overlay histogram showing 293T cells stained with AAA126001 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-MEK1/MAP2K1 Antibody (AAA126001, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)
Figure 1. Western blot analysis of MEK1/MAP2K1 using anti-MEK1/MAP2K1 antibody (AAA126001).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human HeLa whole cell lysates,Lane 2: human Daudi whole cell lysates,Lane 3: rat lung tissue lysates,Lane 4: rat brain tissue lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MEK1/MAP2K1 antigen affinity purified polyclonal antibody (#AAA126001) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for MEK1/MAP2K1 at approximately 45 kDa. The expected band size for MEK1/MAP2K1 is at 45 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-MAP2K1 antibody	Dual specificity mitogen-activated protein kinase kinase 1 is an enzyme that in humans is encoded by the MAP2K1 gene. The protein encoded by this gene is a member of the dual specificity protein kinase family, which acts as a mitogen-activated protein(MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases(ERKs), act as an integration point for multiple biochemical signals. This protein kinase lies upstream of MAP kinases and stimulates the enzymatic activity of MAP kinases upon activation by a wide variety of extra- and intracellular signals. As an essential component of the MAP kinase signal transduction pathway, this kinase is involved in many cellular processes such as proliferation, differentiation, transcription regulation and development. Rampoldi et al.(1997) localized the MAP2K1 gene to 15q22.1-q22.33.
Product Categories/Family for anti-MAP2K1 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	5579478
NCBI GeneID	5604
NCBI Accession #	NP_002746.1
NCBI GenBank Nucleotide #	NM_002755.3
UniProt Accession #	Q02750
Molecular Weight	43,439 Da
NCBI Official Full Name	dual specificity mitogen-activated protein kinase kinase 1
NCBI Official Synonym Full Names	mitogen-activated protein kinase kinase 1
NCBI Official Symbol	MAP2K1
NCBI Official Synonym Symbols	MEK1; MKK1; MAPKK1; PRKMK1

NCBI Protein Information	dual specificity mitogen-activated protein kinase kinase 1; MEK 1; MAPKK 1; MAPK/ERK kinase 1; ERK activator kinase 1; protein kinase, mitogen-activated, kinase 1 (MAP kinase kinase 1)
UniProt Protein Name	Dual specificity mitogen-activated protein kinase kinase 1
UniProt Gene Name	MAP2K1
UniProt Synonym Gene Names	MEK1; PRKMK1; MAP kinase kinase 1; MAPKK 1; MKK1; MEK 1
UniProt Entry Name	MP2K1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mek1-map2k1-antibody-western-blot-flow-cytometry-elisa/126001>