

Rabbit TNF Receptor I/TNFRSF1A Polyclonal Antibody | anti-TNFRSF1A antibody

Anti-TNF Receptor I/TNFRSF1A Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126002	Rabbit	Human, Mouse, Rat	Western Blot, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS

TNF Receptor I/TNFRSF1A; Polyclonal Antibody; Anti-TNF Receptor I/TNFRSF1A Antibody Picoband; anti-TNFRSF1A antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-TNFRSF1A antibody	WB (Western Blot), FCM/FACS (Flow Cytometry), ELISA
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human TNF Receptor I/TNFRSF1A recombinant protein (Position: N336-A440).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)

Figure 2. Flow Cytometry analysis of U937 cells using anti-TNF Receptor I/TNFRSF1A antibody (AAA126002).Overlay histogram showing U937 cells stained with AAA126002 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-TNF Receptor I/TNFRSF1A Antibody (AAA126002, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)

Figure 1. Western blot analysis of TNF Receptor I/TNFRSF1A using anti-TNF Receptor I/TNFRSF1A antibody (AAA126002).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human HepG2 whole cell lysates,Lane 2: human HCCT tissue lysates,Lane 3: human HCCP tissue lysates,Lane 4: human HL-60 whole cell lysates,Lane 5: rat brain tissue lysates,Lane 6: rat C6 whole cell lysates,Lane 7: mouse brain tissue lysates,Lane 8: mouse Neuro-2a whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TNF Receptor I/TNFRSF1A antigen affinity purified polyclonal antibody (#AAA126002) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for TNF Receptor I/TNFRSF1A at approximately 60-70 kDa. The expected band size for TNF Receptor I/TNFRSF1A is at 50 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-TNFRSF1A antibody	Tumor necrosis factor receptor superfamily member 1A (TNFRSF1A), also known as TNFR1, is a protein that in humans is encoded by the TNFRSF1A gene. The protein encoded by this gene is a member of the Tumor necrosis factor receptor superfamily, which also contains TNFRSF1B. The TNFR1 gene is mapped to 12pter-cen. It encodes a protein of 455 amino acids. And this receptor can activate the transcription factor NF-kappaB, mediate apoptosis, and function as a regulator of inflammation.
Product Categories/Family for anti-TNFRSF1A antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507575
NCBI GeneID	7132
NCBI Accession #	NP_001056.1
NCBI GenBank Nucleotide #	NM_001065.3
UniProt Accession #	P19438; Q9UCA4; A8K4X3; B2RDE4; B3KPQ1; B4DQB7; B4E309; D3DUR1
Molecular Weight	50,495 Da
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 1A
NCBI Official Synonym Full Names	tumor necrosis factor receptor superfamily, member 1A
NCBI Official Symbol	TNFRSF1A
NCBI Official Synonym Symbols	FPF; MS5; p55; p60; TBP1; TNF-R; TNFAR; TNFR1; p55-R; CD120a; TNFR55; TNFR60; TNF-R-I; TNF-R55; TNFR1-d2

NCBI Protein Information	tumor necrosis factor receptor superfamily member 1A; TNF-R1; TNF-R1; TNFR-I; tumor necrosis factor-alpha receptor; tumor necrosis factor receptor type 1; tumor necrosis factor binding protein 1; tumor necrosis factor receptor 1A isoform beta
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 1A
UniProt Gene Name	TNFRSF1A
UniProt Synonym Gene Names	TNFAR; TNFR1; TNF-R1; TNF-R1; TNFR-I; TBPI
UniProt Entry Name	TNR1A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/tnf-receptor-i-tnfrsf1a-antibody-western-blot-flow-cytometry-elisa/126002>