

Rabbit anti-Human NOX2/gp91phox/CYBB Polyclonal Antibody | anti-CYBB antibody

Anti-NOX2/gp91phox/CYBB Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126008	Rabbit	Human	ELISA, Flow Cytometry, Functional Assay, Western Blot

PRODUCT SYNONYMS


NOX2/gp91phox/CYBB; Polyclonal Antibody; Anti-NOX2/gp91phox/CYBB Antibody Picoband; anti-CYBB antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-CYBB antibody	ELISA, FCM/FACS (Flow Cytometry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human NOX2/gp91phox/CYBB recombinant protein (Position: F416-H527).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)
Figure 2. Flow Cytometry analysis of U937 cells using anti-NOX2/Gp91phox/CYBB antibody (AAA126008). Overlay histogram showing U937 cells stained with AAA126008 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-NOX2/Gp91phox/CYBB Antibody (AAA126008, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)
Figure 1. Western blot analysis of NOX2/Gp91phox/CYBB using anti-NOX2/Gp91phox/CYBB antibody (AAA126008). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human MCF-7 whole cell lysates, Lane 2: human Raji whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-NOX2/Gp91phox/CYBB antigen affinity purified polyclonal antibody (#AAA126008) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for NOX2/Gp91phox/CYBB at approximately 65 kDa. The expected band size for NOX2/Gp91phox/CYBB is at 65 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-CYBB antibody	NOX2(NADPH OXIDASE 2), also called CYBB(CYTOCHROME b(-245), BETA SUBUNIT), p91-PHOX or GP91-1, is a human gene encoding a glycoprotein. NOX2 is an essential component of phagocytic NADPH-oxidase, a membrane-bound enzyme complex that generates large quantities of microbicidal superoxide and other oxidants upon activation. It is mapped on Xp11.4. NOX2 assembled on DC phagosomes in a gp91-phox subunit-dependent manner, and that reactive oxygen species were produced in a more sustained manner in immature DC phagosomes than in macrophage phagosomes. As a major player in innate immune responses in neutrophils, NOX2 is also involved in adaptive immunity through its activity in DCs. In heart cells, physiologic stretch rapidly activates reduced-form NOX2 to produce reactive oxygen species (ROS) in a process dependent on microtubules (X-ROS signaling).
Product Categories/Family for anti-CYBB antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	115211
NCBI GeneID	1536
UniProt Accession #	P04839; Q2PP16; A8K138
Molecular Weight	65,336 Da
NCBI Official Full Name	Cytochrome b-245 heavy chain
NCBI Official Synonym Full Names	cytochrome b-245, beta polypeptide
NCBI Official Symbol	CYBB
NCBI Official Synonym Symbols	CGD; NOX2; AMCBX2; GP91-1; GP91PHOX; p91-PHOX; GP91-PHOX
NCBI Protein Information	cytochrome b-245 heavy chain; CGD91-phox; NADPH oxidase 2; p22 phagocyte B-cytochrome; cytochrome b558 subunit beta; neutrophil cytochrome b 91 kDa polypeptide; heme-binding membrane glycoprotein gp91phox; superoxide-generating NADPH oxidase heavy chain s

UniProt Protein Name	Cytochrome b-245 heavy chain
UniProt Gene Name	CYBB
UniProt Synonym Gene Names	NOX2; Cytochrome b558 subunit beta
UniProt Entry Name	CY24B_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/nox2-gp91phox-cybb-antibody-flow-cytometry-western-blot-elisa/126008>