

Rabbit ERCC1 Polyclonal Antibody | anti-ERCC1 antibody

Anti-ERCC1 Antibody Picoband

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126016	Rabbit	Human, Mouse, Rat	ELISA, Flow Cytometry, Functional Assay, Immunofluorescence, Immunocytochemistry, Western Blot


PRODUCT SYNONYMS

ERCC1; Polyclonal Antibody; Anti-ERCC1 Antibody Picoband; anti-ERCC1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-ERCC1 antibody	ELISA, FCM/FACS (Flow Cytometry), IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human ERCC1 recombinant protein (Position: R106-E291).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.

IMAGES




FCM/FACS (Flow Cytometry)

Figure 3. Flow Cytometry analysis of HL-60 cells using anti-ERCC1 antibody (AAA126016). Overlay histogram showing HL-60 cells stained with AAA126016 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-ERCC1 Antibody (AAA126016, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IF (Immunofluorescence)

Figure 2. IF analysis of ERCC1 using anti-ERCC1 antibody (AAA126016). ERCC1 was detected in an immunocytochemical section of A549 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-ERCC1 Antibody (AAA126016) overnight at 4 degree C. DyLight594 Conjugated Goat Anti-Rabbit IgG (BA1142) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



WB (Western Blot)

Figure 1. Western blot analysis of ERCC1 using anti-ERCC1 antibody (AAA126016). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human Hela whole cell lysates, Lane 2: human HepG2 whole cell lysates, Lane 3: human COLO-320 whole cell lysates, Lane 4: rat kidney tissue lysates, Lane 5: mouse kidney tissue lysates, Lane 6: mouse NIH/3T3 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ERCC1 antigen affinity purified polyclonal antibody (#AAA126016) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for ERCC1 at approximately 33 kDa. The expected band size for ERCC1 is at 33 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-ERCC1 antibody

DNA excision repair protein ERCC-1 is a protein that in humans is encoded by the ERCC1 gene. The product of this gene functions in the nucleotide excision repair pathway, and is required for the repair of DNA lesions such as those induced by UV light or formed by electrophilic compounds including cisplatin. The encoded protein forms a heterodimer with the XPF endonuclease (also known as ERCC4), and the heterodimeric endonuclease catalyzes the 5' incision in the process of excising the DNA lesion. The heterodimeric endonuclease is also involved in recombinational DNA repair and in the repair of inter-strand crosslinks. Mutations in this gene result in cerebrooculofacioskeletal syndrome, and polymorphisms that alter expression of this gene may play a role in carcinogenesis. Multiple transcript variants encoding different isoforms have been found for this gene. The last exon of this gene overlaps with the CD3e molecule, epsilon associated protein gene on the opposite strand.

Product Categories/Family for anti-ERCC1 antibody

[Primary Antibodies](#), [Rabbit Polyclonal Antibodies](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	260593725
NCBI GeneID	2067
NCBI Accession #	NP_001159521.1
NCBI GenBank Nucleotide #	NM_001166049.1
UniProt Accession #	P07992 ; Q7Z7F5 ; Q96S40
Molecular Weight	32,562 Da
NCBI Official Full Name	DNA excision repair protein ERCC-1 isoform 3
NCBI Official Synonym Full Names	excision repair cross-complementing rodent repair deficiency, complementation group 1 (includes overlapping antisense sequence)
NCBI Official Symbol	ERCC1

NCBI Official Synonym Symbols	UV20; COFS4; RAD10
NCBI Protein Information	DNA excision repair protein ERCC-1
UniProt Protein Name	DNA excision repair protein ERCC-1
UniProt Gene Name	ERCC1
UniProt Entry Name	ERCC1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ercc1-antibody-flow-cytometry-immunofluorescence-immunocytochemistry-western-blot-elisa/126016>