

Rabbit anti-Human ROR gamma/RORC Polyclonal Antibody | anti-RORC antibody

Anti-ROR gamma/RORC Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126026	Rabbit	Human	ELISA, Flow Cytometry, Functional Assay, Western Blot

PRODUCT SYNONYMS


ROR gamma/RORC; Polyclonal Antibody; Anti-ROR gamma/RORC Antibody Picoband; anti-RORC antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-RORC antibody	ELISA, FCM/FACS (Flow Cytometry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human ROR gamma/RORC recombinant protein (Position: Q286-E511).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)
Figure 2. Flow Cytometry analysis of HeLa cells using anti-ROR gamma/RORC antibody (AAA126026).Overlay histogram showing HeLa cells stained with AAA126026 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-ROR gamma/RORC Antibody (AAA126026, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)
Figure 1. Western blot analysis of ROR gamma/RORC using anti-ROR gamma/RORC antibody (AAA126026).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human HCCT tissue lysates,Lane 2: human MOLT-4 whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ROR gamma/RORC antigen affinity purified polyclonal antibody (#AAA126026) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for ROR gamma/RORC at approximately 58 kDa. The expected band size for ROR gamma/RORC is at 58 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-RORC antibody	RAR-related orphan receptor gamma (RORgamma) is a protein that in humans is encoded by the RORC (RAR-related orphan receptor C) gene. The protein encoded by this gene is a DNA-binding transcription factor and is a member of the NR1 subfamily of nuclear hormone receptors. The specific functions of this protein are not known; however, studies of a similar gene in mice have shown that this gene may be essential for lymphoid organogenesis and may play an important regulatory role in thymopoiesis. In addition, studies in mice suggest that the protein encoded by this gene may inhibit the expression of Fas ligand and IL2. Two transcript variants encoding different isoforms have been found for this gene.
Product Categories/Family for anti-RORC antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	48255918
NCBI GeneID	6097
NCBI Accession #	NP_001001523.1
NCBI GenBank Nucleotide #	NM_001001523.1
UniProt Accession #	P51449; Q5SZR9; Q8N5V7; Q8NCY8
Molecular Weight	58,195 Da
NCBI Official Full Name	nuclear receptor ROR-gamma isoform b
NCBI Official Synonym Full Names	RAR-related orphan receptor C
NCBI Official Symbol	RORC
NCBI Official Synonym Symbols	TOR; RORG; RZRG; NR1F3; RZR-GAMMA

NCBI Protein Information	nuclear receptor ROR-gamma; nuclear receptor RZR-gamma; retinoic acid-binding receptor gamma; retinoid-related orphan receptor gamma; RAR-related orphan receptor C, isoform a; RAR-related orphan nuclear receptor variant 2; nuclear receptor subfamily 1 gro
UniProt Protein Name	Nuclear receptor ROR-gamma
UniProt Gene Name	RORC
UniProt Synonym Gene Names	NR1F3; RORG; RZRG
UniProt Entry Name	RORG_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ror-gamma-rorc-antibody-flow-cytometry-western-blot-elisa/126026>