

Rabbit Factor H/CFH Polyclonal Antibody | anti-CFH antibody

Anti-Factor H/CFH Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126039	Rabbit	Human, Mouse, Rat	ELISA, Flow Cytometry, Functional Assay, Western Blot

PRODUCT SYNONYMS


Factor H/CFH; Polyclonal Antibody; Anti-Factor H/CFH Antibody Picoband; anti-CFH antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-CFH antibody	ELISA, FCM/FACS (Flow Cytometry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human Factor H/CFH recombinant protein (Position: N115-A998).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)
Figure 2. Flow Cytometry analysis of U251 cells using anti-Factor H/CFH antibody (AAA126039).Overlay histogram showing U251 cells stained with AAA126039 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Factor H/CFH Antibody (AAA126039, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)
Figure 1. Western blot analysis of Factor H/CFH using anti-Factor H/CFH antibody (AAA126039).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human HCCP tissue lysates,Lane 2: rat liver tissue lysates,Lane 3: mouse liver tissue lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Factor H/CFH antigen affinity purified polyclonal antibody (#AAA126039) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for Factor H/CFH at approximately 150 kDa. The expected band size for Factor H/CFH is at 139 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-CFH antibody	Factor H is a member of the regulators of complement activation family and is a complement control protein. This gene is a member of the Regulator of Complement Activation (RCA) gene cluster and encodes a protein with twenty short consensus repeat (SCR) domains. This protein is secreted into the bloodstream and has an essential role in the regulation of complement activation, restricting this innate defense mechanism to microbial infections. Mutations in this gene have been associated with hemolytic-uremic syndrome (HUS) and chronic hypocomplementemic nephropathy. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.
Product Categories/Family for anti-CFH antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	758073
NCBI GeneID	3075
UniProt Accession #	P08603; P78435; Q14570; Q2TAZ5; Q38G77; Q5TFM3; Q8N708; Q9NU86; A5PL14
Molecular Weight	139,096 Da
NCBI Official Full Name	complement factor H
NCBI Official Synonym Full Names	complement factor H
NCBI Official Symbol	CFH
NCBI Official Synonym Symbols	FH; HF; HF1; HF2; HUS; FHL1; AHUS1; AMBP1; ARMD4; ARMS1; CFHL3
NCBI Protein Information	complement factor H; beta-1H; factor H-like 1; beta-1-H-globulin; H factor 1 (complement); H factor 2 (complement); adrenomedullin binding protein; complement factor H, isoform b; age-related maculopathy susceptibility 1
UniProt Protein Name	Complement factor H
UniProt Gene Name	CFH

UniProt Synonym Gene Names	HF; HF1; HF2
UniProt Entry Name	CFAH_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/factor-h-cfh-antibody-flow-cytometry-western-blot-elisa/126039>