

Rabbit anti-Human CD2 Polyclonal Antibody | anti-CD2 antibody

Anti-CD2 Antibody Picoband


CATALOG # AAA126043	HOST Rabbit	REACTIVITY Human	APPLICATIONS ELISA, Flow Cytometry, Functional Assay, Western Blot
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS


CD2; Polyclonal Antibody; Anti-CD2 Antibody Picoband; anti-CD2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-CD2 antibody	ELISA, FCM/FACS (Flow Cytometry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human CD2 recombinant protein (Position: V23-D209).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)
Figure 2. Flow Cytometry analysis of JK cells using anti-CD2 antibody (AAA126043).Overlay histogram showing JK cells stained with AAA126043 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-CD2 Antibody (AAA126043, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. Dylight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)
Figure 1. Western blot analysis of CD2 using anti-CD2 antibody (AAA126043).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human MOLT-4 whole cell lysates,Lane 2: human Jurkat whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CD2 antigen affinity purified polyclonal antibody (#AAA126043) at 0.25 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for CD2 at approximately 50 kDa. The expected band size for CD2 is at 39 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-CD2 antibody	CD2 (cluster of differentiation 2) is a cell adhesion molecule found on the surface of T cells and natural killer (NK) cells. It has also been called T-cell surface antigen T11/Leu-5, LFA-2, LFA-3 receptor, erythrocyte receptor and rosette receptor. Monoclonal antibodies directed against CD2 inhibit the formation of rosettes with sheep erythrocytes, indicating that CD2 is the erythrocyte receptor or is closely associated with it. CD2 is one of the earliest T-cell markers, being present on more than 95% of thymocytes. Due to its structural characteristics, CD2 is a member of the immunoglobulin superfamily; it possesses two immunoglobulin-like domains in its extracellular portion. The localization of CD2 to 1p13 is established by in situ hybridization. CD2 interacts with other adhesion molecules, such as lymphocyte function-associated antigen-3 (LFA-3/CD58) in humans, or CD48 in rodents, which are expressed on the surfaces of other cells. With the use of transgenic mice, such an LCR is identified within the 3-prime flanking region of the human CD2 gene.
Product Categories/Family for anti-CD2 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	156071472
NCBI GeneID	914
NCBI Accession #	NP_001758.2
NCBI GenBank Nucleotide #	NM_001767.3
UniProt Accession #	P06729; Q96TE5
Molecular Weight	39,448 Da
NCBI Official Full Name	T-cell surface antigen CD2
NCBI Official Synonym Full Names	CD2 molecule
NCBI Official Symbol	CD2

NCBI Official Synonym Symbols	T11; SRBC; LFA-2
NCBI Protein Information	T-cell surface antigen CD2; CD2 antigen (p50), sheep red blood cell receptor; LFA-3 receptor; T-cell surface antigen T11/Leu-5; erythrocyte receptor; lymphocyte-function antigen-2; rosette receptor
UniProt Protein Name	T-cell surface antigen CD2
UniProt Gene Name	CD2
UniProt Synonym Gene Names	SRBC
UniProt Entry Name	CD2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd2-antibody-flow-cytometry-western-blot-elisa/126043>