

Rabbit anti-Human, Rat L1CAM Polyclonal Antibody | anti-L1CAM antibody

Anti-L1CAM Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126066	Rabbit	Human, Rat	Western Blot, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS


L1CAM; Polyclonal Antibody; Anti-L1CAM Antibody Picoband; anti-L1CAM antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-L1CAM antibody	WB (Western Blot), FCM/FACS (Flow Cytometry), ELISA
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human L1CAM recombinant protein (Position: E223-Q1218).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)
Figure 2. Flow Cytometry analysis of Caco-2 cells using anti-L1CAM antibody (AAA126066).Overlay histogram showing Caco-2 cells stained with AAA126066 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-L1CAM Antibody (AAA126066, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)
Figure 1. Western blot analysis of L1CAM using anti-L1CAM antibody (AAA126066).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human Hela whole cell lysates,Lane 2: rat brain tissue lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-L1CAM antigen affinity purified polyclonal antibody (#AAA126066) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for L1CAM at approximately 220 kDa. The expected band size for L1CAM is at 140 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-L1CAM antibody	L1, also known as L1CAM, is a transmembrane protein member of the L1 protein family, encoded by the L1CAM gene. The protein encoded by this gene is an axonal glycoprotein belonging to the immunoglobulin supergene family. The ectodomain, consisting of several immunoglobulin-like domains and fibronectin-like repeats (type III), is linked via a single transmembrane sequence to a conserved cytoplasmic domain. This cell adhesion molecule plays an important role in nervous system development, including neuronal migration and differentiation. Mutations in the gene cause X-linked neurological syndromes known as CRASH (corpus callosum hypoplasia, retardation, aphasia, spastic paraplegia and hydrocephalus). Alternative splicing of this gene results in multiple transcript variants, some of which include an alternate exon that is considered to be specific to neurons.
Product Categories/Family for anti-L1CAM antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557707
NCBI GeneID	3897
NCBI Accession #	NP_000416.1
NCBI GenBank Nucleotide #	NM_000425.4
UniProt Accession #	P32004; Q8TA87; A0AV65; A4ZYW4; B2RMU7; G3XAF4
Molecular Weight	140,003 Da
NCBI Official Full Name	neural cell adhesion molecule L1 isoform 1
NCBI Official Synonym Full Names	L1 cell adhesion molecule
NCBI Official Symbol	L1CAM
NCBI Official Synonym Symbols	S10; HSAS; MASA; MIC5; SPG1; CAML1; CD171; HSAS1; N-CAML1; NCAM-L1; N-CAM-L1

NCBI Protein Information	neural cell adhesion molecule L1; antigen identified by monoclonal antibody R1
UniProt Protein Name	Neural cell adhesion molecule L1
UniProt Gene Name	L1CAM
UniProt Synonym Gene Names	CAML1; MIC5; N-CAM-L1; NCAM-L1
UniProt Entry Name	L1CAM_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/l1cam-antibody-western-blot-flow-cytometry-elisa/126066>