

Rabbit BUB1 Polyclonal Antibody | anti-BUB1 antibody

Anti-BUB1 Antibody Picoband

CATALOG # AAA126073	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS ELISA, Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

BUB1; Polyclonal Antibody; Anti-BUB1 Antibody Picoband; anti-BUB1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-BUB1 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human BUB1 recombinant protein (Position: E26-K1085).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.

IMAGES




IHC (Immunohistochemistry)

Figure 4. IHC analysis of BUB1 using anti-BUB1 antibody (AAA126073). BUB1 was detected in a paraffin-embedded section of rat thymus tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-BUB1 Antibody (AAA126073) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 3. IHC analysis of BUB1 using anti-BUB1 antibody (AAA126073). BUB1 was detected in a paraffin-embedded section of mouse thymus tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-BUB1 Antibody (AAA126073) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of BUB1 using anti-BUB1 antibody (AAA126073). BUB1 was detected in a paraffin-embedded section of human bladder cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-BUB1 Antibody (AAA126073) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of BUB1 using anti-BUB1 antibody (AAA126073). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human Hela whole cell lysates, Lane 2: human 293T whole cell lysates, Lane 3: human K562 whole cell lysates, Lane 4: mouse testis tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-BUB1 antigen affinity purified polyclonal antibody (#AAA126073) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for BUB1 at approximately 130 kDa. The expected band size for BUB1 is at 122 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-BUB1 antibody

BUB1, also known as mitotic checkpoint serine/threonine kinase, is an enzyme that in humans is encoded by the BUB1 gene. It is mapped to 2q13. BUB1 is first identified in genetic screens of *Saccharomyces cerevisiae*. The protein is bound to kinetochores and plays a key role in the establishment of the mitotic spindle checkpoint and chromosome congression. The mitotic checkpoint kinase is evolutionary conserved in organisms as diverse as *Saccharomyces cerevisiae* and humans. Loss-of-function mutations or absence of BUB1 has been reported to result in aneuploidy, chromosomal instability (CIN) and premature senescence. The protein kinase BUB1 possesses versatile and distinct functions during the cell cycle, mainly in the SAC and chromosome alignment during metaphase.

Product Categories/Family for anti-BUB1 antibody

Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4757878
NCBI GeneID	699
NCBI Accession #	NP_004327.1
NCBI GenBank Nucleotide #	NM_004336.3
UniProt Accession #	O43683; O43430; O43643; O60626; Q53QE4
Molecular Weight	122,375 Da
NCBI Official Full Name	mitotic checkpoint serine/threonine-protein kinase BUB1
NCBI Official Synonym Full Names	BUB1 mitotic checkpoint serine/threonine kinase
NCBI Official Symbol	BUB1
NCBI Official Synonym Symbols	BUB1A; BUB1L; hBUB1

NCBI Protein Information	mitotic checkpoint serine/threonine-protein kinase BUB1; mitotic spindle checkpoint kinase; putative serine/threonine-protein kinase; budding uninhibited by benzimidazoles 1 homolog; BUB1 budding uninhibited by benzimidazoles 1 homolog
UniProt Protein Name	Mitotic checkpoint serine/threonine-protein kinase BUB1
UniProt Gene Name	BUB1
UniProt Synonym Gene Names	BUB1L; hBUB1
UniProt Entry Name	BUB1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/bub1-antibody-immunohistochemistry-western-blot-elisa/126073>