

Rabbit anti-Human GSK3 beta/GSK3B Polyclonal Antibody | anti-GSK3B antibody

Anti-GSK3 beta/GSK3B Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126076	Rabbit	Human	ELISA, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

GSK3 beta/GSK3B; Polyclonal Antibody; Anti-GSK3 beta/GSK3B Antibody Picoband; anti-GSK3B antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-GSK3B antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human GSK3 beta/GSK3B recombinant protein (Position: M1-T420).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




IHC (Immunohistochemistry)

Figure 3. IHC analysis of GSK3 beta/GSK3B using anti-GSK3 beta/GSK3B antibody (AAA126076).GSK3 beta/GSK3B was detected in a paraffin-embedded section of human ovarian cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-GSK3 beta/GSK3B Antibody (AAA126076) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of GSK3 beta/GSK3B using anti-GSK3 beta/GSK3B antibody (AAA126076).GSK3 beta/GSK3B was detected in a paraffin-embedded section of human lung adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-GSK3 beta/GSK3B Antibody (AAA126076) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of GSK3 beta/GSK3B using anti-GSK3 beta/GSK3B antibody (AAA126076).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human U-87MG whole cell lysates,Lane 2: human HeLa whole cell lysates,Lane 3: human A431 whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-GSK3 beta/GSK3B antigen affinity purified polyclonal antibody (#AAA126076) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for GSK3 beta/GSK3B at approximately 46 kDa. The expected band size for GSK3 beta/GSK3B is at 46 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-GSK3B antibody	Glycogen synthase kinase-3 beta, (GSK-3 beta), is an enzyme that in humans is encoded by the GSK3B gene. The protein encoded by this gene is a serine-threonine kinase belonging to the glycogen synthase kinase subfamily. It is a negative regulator of glucose homeostasis and is involved in energy metabolism, inflammation, ER-stress, mitochondrial dysfunction, and apoptotic pathways. Defects in this gene have been associated with Parkinson disease and Alzheimer disease.
Product Categories/Family for anti-GSK3B antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	225903437
NCBI GeneID	2932
NCBI Accession #	NP_001139628.1
NCBI GenBank Nucleotide #	NM_001146156.1
UniProt Accession #	P49841; Q9BWH3; Q9UL47; D3DN89
Molecular Weight	46,744 Da
NCBI Official Full Name	glycogen synthase kinase-3 beta isoform 2
NCBI Official Synonym Full Names	glycogen synthase kinase 3 beta
NCBI Official Symbol	GSK3B
NCBI Protein Information	glycogen synthase kinase-3 beta; GSK-3 beta; GSK3beta isoform; serine/threonine-protein kinase GSK3B
UniProt Protein Name	Glycogen synthase kinase-3 beta
UniProt Gene Name	GSK3B
UniProt Synonym Gene Names	GSK-3 beta
UniProt Entry Name	GSK3B_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/gsk3-beta-gsk3b-antibody-immunohistochemistry-western-blot-elisa/126076>