

Rabbit anti-Human, Mouse MX1 Polyclonal Antibody | anti-MX1 antibody

Anti-MX1 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126088	Rabbit	Human, Mouse	ELISA, Flow Cytometry, Functional Assay, Western Blot

PRODUCT SYNONYMS


MX1; Polyclonal Antibody; Anti-MX1 Antibody Picoband; anti-MX1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-MX1 antibody	ELISA, FCM/FACS (Flow Cytometry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human MX1 recombinant protein (Position: N116-H595).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)
Figure 2. Flow Cytometry analysis of RT4 cells using anti-MX1 antibody (AAA126088).Overlay histogram showing RT4 cells stained with AAA126088 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-MX1 Antibody (AAA126088, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)
Figure 1. Western blot analysis of MX1 using anti-MX1 antibody (AAA126088).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human Hela whole cell lysates,Lane 2: human A549 whole cell lysates,Lane 3: mouse lung tissue lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MX1 antigen affinity purified polyclonal antibody (#AAA126088) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for MX1 at approximately 76 kDa. The expected band size for MX1 is at 76 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-MX1 antibody	Interferon-induced GTP-binding protein Mx1 is a protein that in humans is encoded by the MX1 gene. This gene encodes a guanosine triphosphate (GTP)-metabolizing protein that participates in the cellular antiviral response. The encoded protein is induced by type I and type II interferons and antagonizes the replication process of several different RNA and DNA viruses. There is a related gene located adjacent to this gene on chromosome 21, and there are multiple pseudogenes located in a cluster on chromosome 4. Alternative splicing results in multiple transcript variants.
Product Categories/Family for anti-MX1 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	251757499
NCBI GeneID	4599
UniProt Accession #	P20591; Q86YP5; Q96CI3; B2RDA5; B3KU10; C9IYV7; C9J8D6; C9JN19; C9JN88; C9JUL1; C9JZS6; D3DSI8
Molecular Weight	662
NCBI Official Full Name	Interferon-induced GTP-binding protein Mx1
NCBI Official Synonym Full Names	MX dynamin-like GTPase 1
NCBI Official Symbol	MX1
NCBI Official Synonym Symbols	MX; MxA; IFI78; IFI-78K
NCBI Protein Information	interferon-induced GTP-binding protein Mx1; myxoma resistance protein 1; interferon-induced protein p78; interferon-inducible protein p78; interferon-regulated resistance GTP-binding protein MxA; myxovirus (influenza virus) resistance 1, interferon-induci
UniProt Protein Name	Interferon-induced GTP-binding protein Mx1

UniProt Gene Name	MX1
UniProt Synonym Gene Names	IFI-78K
UniProt Entry Name	MX1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mx1-antibody-flow-cytometry-western-blot-elisa/126088>