

Rabbit anti-Human MRP1/ABCC1 Polyclonal Antibody | anti-ABCC1 antibody

Anti-MRP1/ABCC1 Antibody Picoband


CATALOG # AAA126089	HOST Rabbit	REACTIVITY Human	APPLICATIONS ELISA, Flow Cytometry, Functional Assay, Western Blot
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS


MRP1/ABCC1; Polyclonal Antibody; Anti-MRP1/ABCC1 Antibody Picoband; anti-ABCC1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-ABCC1 antibody	ELISA, FCM/FACS (Flow Cytometry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human MRP1/ABCC1 recombinant protein (Position: S239-K357).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)
Figure 2. Flow Cytometry analysis of SiHa cells using anti-MRP1/ABCC1 antibody (AAA126089).Overlay histogram showing SiHa cells stained with AAA126089 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-MRP1/ABCC1 Antibody (AAA126089, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)
Figure 1. Western blot analysis of MRP1/ABCC1 using anti-MRP1/ABCC1 antibody (AAA126089).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human PANC-1 whole cell lysates,Lane 2: human A549 whole cell lysates,Lane 3: human HepG2 whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MRP1/ABCC1 antigen affinity purified polyclonal antibody (#AAA126089) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for MRP1/ABCC1 at approximately 220 kDa. The expected band size for MRP1/ABCC1 is at 220 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-ABCC1 antibody	Multidrug resistance-associated protein 1 (MRP1) is a protein that in humans is encoded by the ABCC1 gene. The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra-and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This full transporter is a member of the MRP subfamily which is involved in multi-drug resistance. This protein functions as a multispecific organic anion transporter, with oxidized glutathione, cysteinyl leukotrienes, and activated aflatoxin B1 as substrates. This protein also transports glucuronides and sulfate conjugates of steroid hormones and bile salts. Alternatively spliced variants of this gene have been described but their full-length nature is unknown.
Product Categories/Family for anti-ABCC1 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	296439301
NCBI GeneID	4363
UniProt Accession #	P33527; O14819; O43333; P78419; Q59GI9; Q9UQ97; Q9UQ99; Q9UQA0; A3RJX2; C9JPJ4
Molecular Weight	1531
NCBI Official Full Name	Multidrug resistance-associated protein 1
NCBI Official Synonym Full Names	ATP-binding cassette, sub-family C (CFTR/MRP), member 1
NCBI Official Symbol	ABCC1
NCBI Official Synonym Symbols	MRP; ABCC; GS-X; MRP1; ABC29
NCBI Protein Information	multidrug resistance-associated protein 1; LTC4 transporter; leukotriene C(4) transporter; ATP-binding cassette transporter variant ABCC1delta-ex13; ATP-binding cassette transporter variant ABCC1delta-ex25; ATP-binding cassette transporter variant ABCC1de

UniProt Protein Name	Multidrug resistance-associated protein 1
UniProt Gene Name	ABCC1
UniProt Synonym Gene Names	MRP; MRP1; LTC4 transporter
UniProt Entry Name	MRP1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mrp1-abcc1-antibody-flow-cytometry-western-blot-elisa/126089>