

Rabbit ASK1/MAP3K5 Polyclonal Antibody | anti-MAP3K5 antibody

Anti-ASK1/MAP3K5 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126095	Rabbit	Human, Mouse, Rat	Western Blot, Immunocytochemistry, Immunohistochemistry, Immunofluorescence, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS

ASK1/MAP3K5; Polyclonal Antibody; Anti-ASK1/MAP3K5 Antibody Picoband; anti-MAP3K5 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-MAP3K5 antibody	WB (Western Blot), ICC (Immunocytochemistry), IHC (Immunohistochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry), ELISA
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human ASK1/MAP3K5 recombinant protein (Position: A178-K1112).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




FCM/FACS (Flow Cytometry)

Figure 5. Flow Cytometry analysis of Raji cells using anti-ASK1/MAP3K5 antibody (AAA126095).Overlay histogram showing Raji cells stained with AAA126095 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-ASK1/MAP3K5 Antibody (AAA126095, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




IHC (Immunohistochemisry)

Figure 3. IHC analysis of ASK1/MAP3K5 using anti-ASK1/MAP3K5 antibody (AAA126095).ASK1/MAP3K5 was detected in a paraffin-embedded section of human spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-ASK1/MAP3K5 Antibody (AAA126095) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IF (Immunofluorescence)

Figure 4. IF analysis of ASK1/MAP3K5 using anti-ASK1/MAP3K5 antibody (AAA126095).ASK1/MAP3K5 was detected in an immunocytochemical section of MCF-7 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-ASK1/MAP3K5 Antibody (AAA126095) overnight at 4 degree C. DyLight488 Conjugated Goat Anti-Rabbit IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



IHC (Immunohiostchemistry)

Figure 2. IHC analysis of ASK1/MAP3K5 using anti-ASK1/MAP3K5 antibody (AAA126095).ASK1/MAP3K5 was detected in a paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-ASK1/MAP3K5 Antibody (AAA126095) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of ASK1/MAP3K5 using anti-ASK1/MAP3K5 antibody (AAA126095). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human RT4 whole cell lysates, Lane 2: human HEL whole cell lysates, Lane 3: human HL-60 whole cell lysates, Lane 4: human THP-1 whole cell lysates, Lane 5: rat liver tissue lysates, Lane 6: mouse liver tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ASK1/MAP3K5 antigen affinity purified polyclonal antibody (#AAA126095) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for ASK1/MAP3K5 at approximately 155 kDa. The expected band size for ASK1/MAP3K5 is at 155 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-MAP3K5 antibody

Apoptosis signal-regulating kinase 1 (ASK1) also known as mitogen-activated protein kinase 5 (MAP3K5) is a member of MAP kinase family and as such a part of mitogen-activated protein kinase pathway. Mitogen-activated protein kinase (MAPK) signaling cascades include MAPK or extracellular signal-regulated kinase (ERK), MAPK kinase (MKK or MEK), and MAPK kinase kinase (MAPKKK or MEKK). MAPKK kinase/MEKK phosphorylates and activates its downstream protein kinase, MAPK kinase/MEK, which in turn activates MAPK. The kinases of these signaling cascades are highly conserved, and homologs exist in yeast, Drosophila, and mammalian cells. MAPKKK5 contains 1,374 amino acids with all 11 kinase subdomains. Northern blot analysis shows that MAPKKK5 transcript is abundantly expressed in human heart and pancreas. The MAPKKK5 protein phosphorylates and activates MKK4 (aliases SERK1, MAPKK4) in vitro, and activates c-Jun N-terminal kinase (JNK)/stress-activated protein kinase (SAPK) during transient expression in COS and 293 cells; MAPKKK5 does not activate MAPK/ERK.

Product Categories/Family for anti-MAP3K5 antibody

[Primary Antibodies](#), [Rabbit Polyclonal Antibodies](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	5174547
NCBI GeneID	4217
NCBI Accession #	NP_005914.1
NCBI GenBank Nucleotide #	NM_005923.3
UniProt Accession #	Q99683 ; Q5THN3 ; Q99461
Molecular Weight	154,537 Da
NCBI Official Full Name	mitogen-activated protein kinase kinase kinase 5
NCBI Official Synonym Full Names	mitogen-activated protein kinase kinase kinase 5
NCBI Official Symbol	MAP3K5
NCBI Official Synonym Symbols	ASK1; MEKK5; MAPKKK5

NCBI Protein Information	mitogen-activated protein kinase kinase kinase 5; ASK-1; MEKK 5; MEK kinase 5; MAP/ERK kinase kinase 5; MAPK/ERK kinase kinase 5; apoptosis signal regulating kinase 1; apoptosis signal-regulating kinase 1
UniProt Protein Name	Mitogen-activated protein kinase kinase kinase 5
UniProt Gene Name	MAP3K5
UniProt Synonym Gene Names	ASK1; MAPKKK5; MEKK5; ASK-1; MEK kinase 5; MEKK 5
UniProt Entry Name	M3K5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ask1-map3k5-antibody-western-blot-immunocytochemistry-immunohistochemistry-immunofluorescence-flow-cytometry-elisa/126095>