

Rabbit anti-Human BST2/Tetherin Polyclonal Antibody | anti-BST2 antibody

Anti-BST2/Tetherin Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126096	Rabbit	Human	ELISA, Flow Cytometry, Functional Assay, Western Blot

PRODUCT SYNONYMS


BST2/Tetherin; Polyclonal Antibody; Anti-BST2/Tetherin Antibody Picoband; anti-BST2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-BST2 antibody	ELISA, FCM/FACS (Flow Cytometry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human BST2/Tetherin recombinant protein (Position: N49-S161).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM/FACS (Flow Cytometry)
Figure 2. Flow Cytometry analysis of Raji cells using anti-BST2/Tetherin antibody (AAA126096).Overlay histogram showing Raji cells stained with AAA126096 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-BST2/Tetherin Antibody (AAA126096, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)
Figure 1. Western blot analysis of BST2/Tetherin using anti-BST2/Tetherin antibody (AAA126096).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human HeLa whole cell lysates,Lane 2: human HepG2 whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-BST2/Tetherin antigen affinity purified polyclonal antibody (#AAA126096) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for BST2/Tetherin at approximately 25-45 kDa. The expected band size for BST2/Tetherin is at 20 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-BST2 antibody	Tetherin, also known as bone marrow stromal antigen 2, is a lipid raft associated protein that in humans is encoded by the BST2 gene. Bone marrow stromal cells are involved in the growth and development of B-cells. The specific function of the protein encoded by the bone marrow stromal cell antigen 2 is undetermined; however, this protein may play a role in pre-B-cell growth and in rheumatoid arthritis.
Product Categories/Family for anti-BST2 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4757876
NCBI GeneID	684
NCBI Accession #	NP_004326.1
NCBI GenBank Nucleotide #	NM_004335.2
UniProt Accession #	Q10589; Q53G07; A8K4Y4
Molecular Weight	19,769 Da
NCBI Official Full Name	bone marrow stromal antigen 2
NCBI Official Synonym Full Names	bone marrow stromal cell antigen 2
NCBI Official Symbol	BST2
NCBI Official Synonym Symbols	CD317; TETHERIN
NCBI Protein Information	bone marrow stromal antigen 2; BST-2; NPC-A-7; HM1.24 antigen
UniProt Protein Name	Bone marrow stromal antigen 2

UniProt Gene Name	BST2
UniProt Synonym Gene Names	BST-2
UniProt Entry Name	BST2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/bst2-tetherin-antibody-flow-cytometry-western-blot-elisa/126096>