

Rabbit EIF5A Polyclonal Antibody | anti-EIF5A antibody

Anti-EIF5A Antibody Picoband

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126168	Rabbit	Human, Mouse, Rat	ELISA, Flow Cytometry, Functional Assay, Immunofluorescence, Immunocytochemistry, Western Blot


PRODUCT SYNONYMS

EIF5A; Polyclonal Antibody; Anti-EIF5A Antibody Picoband; anti-EIF5A antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-EIF5A antibody	ELISA, FCM/FACS (Flow Cytometry), IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human EIF5A recombinant protein (Position: E107-K154).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.

IMAGES




FCM/FACS (Flow Cytometry)

Figure 4. Flow Cytometry analysis of PC-3 cells using anti-EIF5A antibody (AAA126168). Overlay histogram showing PC-3 cells stained with AAA126168 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-EIF5A Antibody (AAA126168, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




FCM/FACS (Flow Cytometry)

Figure 3. Flow Cytometry analysis of Neuro-2a cells using anti-EIF5A antibody (AAA126168). Overlay histogram showing Neuro-2a cells stained with AAA126168 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-EIF5A Antibody (AAA126168, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IF (Immunofluorescence)

Figure 2. IF analysis of EIF5A using anti-EIF5A antibody (AAA126168). EIF5A was detected in an immunocytochemical section of A549 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-EIF5A Antibody (AAA126168) overnight at 4 degree C. DyLight488 Conjugated Goat Anti-Rabbit IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



WB (Western Blot)

Figure 1. Western blot analysis of EIF5A using anti-EIF5A antibody (AAA126168). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human Hela whole cell lysates, Lane 2: human Jurkat whole cell lysates, Lane 3: rat kidney tissue lysates, Lane 4: mouse kidney tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-EIF5A antigen affinity purified polyclonal antibody (#AAA126168) at 0.25 ug/mL overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for EIF5A at approximately 18 kDa. The expected band size for EIF5A is at 18 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-EIF5A antibody

Eukaryotic translation initiation factor 5A-1 is a protein that in humans is encoded by the EIF5A gene. Eukaryotic initiation factor 5A (eIF5A) is an mRNA-binding protein that is involved in translation elongation and plays an important role in promoting translation of polyproline motifs. The eIF5A (eIF5A1) and eIF5A2 genes encode the two vertebrate eIF5A isoforms. While eIF5A1 is expressed constitutively in all tissues, eIF5A2 is mainly expressed in gonads. eIF5A and eIF5A2 are the only identified proteins that contain the distinctive amino acid hypusine, which is generated posttranslationally from lysine through a highly conserved polyamine metabolism pathway. eIF5A function and hypusine modification are both essential for cell proliferation, as knock down of eIF5A expression or blocking eIF5A hypusine modification suppresses cancer cell proliferation. Interestingly, eIF5A is an identified component of a tumor suppressor network of the polyamine-hypusine axis. Co-suppression of both eIF5A and adenosylmethionine decarboxylase 1 (AMD1) promotes lymphomagenesis in mice, while heterozygous deletions of the corresponding AMD1 and eIF5A genes often occur together in human lymphomas.

Product Categories/Family for anti-EIF5A antibody

[Primary Antibodies](#), [Rabbit Polyclonal Antibodies](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	54037409
NCBI GeneID	1984
UniProt Accession #	P63241; P10159; Q16182; Q7L7L3; Q7Z4L1; Q9D0G2; A8K9A0; D3DTP2; AAB29229
Molecular Weight	154
NCBI Official Full Name	Eukaryotic translation initiation factor 5A-1
NCBI Official Synonym Full Names	eukaryotic translation initiation factor 5A
NCBI Official Symbol	EIF5A
NCBI Official Synonym Symbols	EIF-5A; EIF5A1; eIF5AI

NCBI Protein Information	eukaryotic translation initiation factor 5A-1; eIF-4D; eIF-5A1; eIF-5A-1; rev-binding factor; eukaryotic initiation factor 5A
UniProt Protein Name	Eukaryotic translation initiation factor 5A-1
UniProt Gene Name	EIF5A
UniProt Synonym Gene Names	eIF-5A-1; eIF-5A1; eIF-5A
UniProt Entry Name	IF5A1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/eif5a-antibody-flow-cytometry-immunofluorescence-immunocytochemistry-western-blot-elisa/126168>