

Rabbit RED1/ADARB1 Polyclonal Antibody | anti-ADARB1 antibody

Anti-RED1/ADARB1 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126178	Rabbit	Human, Mouse, Rat	ELISA, Flow Cytometry, Functional Assay, Western Blot

PRODUCT SYNONYMS

RED1/ADARB1; Polyclonal Antibody; Anti-RED1/ADARB1 Antibody Picoband; anti-ADARB1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Natural and recombinant Mouse Col18a1
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-ADARB1 antibody	ELISA, FCM/FACS (Flow Cytometry), WB (Western Blot)
Assay Type	Sandwich
Samples	Cell culture supernatants, cell lysates, serum and plasma (heparin, EDTA)
Detection Range	156 pg/ml - 10,000 pg/ml
Sensitivity	<10 pg/ml
Intra-assay Precision	Intra-Assay Precision (Precision within an assay): Three samples of known concentration were tested on one plate to assess intra-assay precision.
Inter-assay Precision	Inter-Assay Precision (Precision across assays): Three samples of known concentration were tested in separate assays to assess inter-assay precision.
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




FCM/FACS (Flow Cytometry)

Figure 3. Flow Cytometry analysis of HEL cells using anti-RED1/ADARB1 antibody (AAA126178).Overlay histogram showing HEL cells stained with AAA126178 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-RED1/ADARB1 Antibody (AAA126178, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



FCM/FACS (Flow Cytometry)

Figure 2. Flow Cytometry analysis of 293T cells using anti-RED1/ADARB1 antibody (AAA126178).Overlay histogram showing 293T cells stained with AAA126178 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-RED1/ADARB1 Antibody (AAA126178, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



WB (Western Blot)

Figure 1. Western blot analysis of RED1/ADARB1 using anti-RED1/ADARB1 antibody (AAA126178).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.Lane 1: human SiHa whole cell lysates,Lane 2: human Hela whole cell lysates,Lane 3: human HEL whole cell lysates,Lane 4: rat brain tissue lysates,Lane 5: mouse brain tissue lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-RED1/ADARB1 antigen affinity purified polyclonal antibody (#AAA126178) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for RED1/ADARB1 at approximately 90 kDa. The expected band size for RED1/ADARB1 is at 81 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-ADARB1 antibody	Background: Endostatin is a naturally-occurring 20-kDa C-terminal fragment derived from type XVIII collagen. It is reported to serve as an anti-angiogenic agent, similar to angiostatin and thrombospondin. And It is produced by proteolytic cleavage of collagen XVIII, a member of the multiplexin family that is characterized by interruptions in the triple helix creating multiple domains, by proteases such as cathepsins. Using a genomic clone as a probe for fluorescence in situ hybridization, Endostatin was mapped the COL18A1 gene to 21q22.3. By immunoprecipitation analysis using membrane fractions of human mammary epithelial cells, It showed that endostatin specifically bound to cell surface nucleolin with high affinity. Blockage of nucleolin with neutralizing antibody or knockdown of nucleolin by RNA interference countered the antiendothelial activity of endostatin and abrogated its antiangiogenic and antitumor activity in vivo. Principle of the Assay: The Quick Picokine™ Mouse Col18a1 Pre-Coated ELISA (Enzyme-Linked Immunosorbent Assay) kit is a solid phase immunoassay specially designed to measure Mouse Col18a1 in cell culture supernatants, cell lysates, serum and plasma (heparin, EDTA). It uses our proprietary Quick ELISA technology. Quick ELISA technology employs capture antibodies conjugated to an affinity tag that is recognized by the polyclonal antibody used to coat our Quick ELISA plates. This approach to sandwich ELISA allows the formation of the antibody-analyte sandwich complex in a single step, significantly reducing assay time. The kit contains recombinant Mouse Col18a1 with immunogen: Expression system for standard: NS0; Immunogen sequence: H1591-K1774. To measure Mouse Col18a1, add standards and samples to the wells, then add antibody cocktail. Wash the wells with PBS or TBS buffer, and add TMB. TMB is an HRP substrate and will be catalyzed to produce a blue color product, which changes into yellow after adding the acidic stop solution. The absorbance of the yellow product is linearly proportional to Mouse Col18a1 in the sample. Read the absorbance of the yellow product in each well using a plate reader, and benchmark the sample wells' readings against the standard curve to determine the concentration of Mouse Col18a1 in the sample.
Product Categories/Family for anti-ADARB1 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4501919
NCBI GeneID	104
NCBI Accession #	NP_001103.1
NCBI GenBank Nucleotide #	NM_001112.3
UniProt Accession #	P78563; O00395; O00465; O00691; O00692; P78555; Q4AE79; Q8NFD1; A6NFK8; A6NJ84; C3TTQ1
Molecular Weight	80,763 Da
NCBI Official Full Name	double-stranded RNA-specific editase 1 isoform 1
NCBI Official Synonym Full Names	adenosine deaminase, RNA-specific, B1
NCBI Official Symbol	ADARB1
NCBI Official Synonym Symbols	RED1; ADAR2; DRABA2; DRADA2
NCBI Protein Information	double-stranded RNA-specific editase 1; RNA editase; RED1 homolog; RNA-editing enzyme 1; RNA editing deaminase 1; RNA-editing deaminase 1; dsRNA adenosine deaminase DRADA2; adenosine deaminase, RNA-specific, B1 (RED1 homolog rat); adenosine deaminase, RNA
UniProt Protein Name	Double-stranded RNA-specific editase 1
UniProt Gene Name	ADARB1
UniProt Synonym Gene Names	ADAR2; DRADA2; RED1
UniProt Entry Name	RED1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

