

Rabbit anti-Human SCA2/ATXN2 Polyclonal Antibody | anti-ATXN2 antibody

Anti-SCA2/ATXN2 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126189	Rabbit	Human	ELISA, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS


SCA2/ATXN2; Polyclonal Antibody; Anti-SCA2/ATXN2 Antibody Picoband; anti-ATXN2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-ATXN2 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human SCA2/ATXN2 recombinant protein (Position: Q1278-L1313).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




IHC (Immunohistochemistry)
Figure 5. IHC analysis of SCA2/ATXN2 using anti-SCA2/ATXN2 antibody (AAA126189).SCA2/ATXN2 was detected in a paraffin-embedded section of human breast cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-SCA2/ATXN2 Antibody (AAA126189) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)
Figure 4. IHC analysis of SCA2/ATXN2 using anti-SCA2/ATXN2 antibody (AAA126189).SCA2/ATXN2 was detected in a paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-SCA2/ATXN2 Antibody (AAA126189) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)
Figure 3. IHC analysis of SCA2/ATXN2 using anti-SCA2/ATXN2 antibody (AAA126189).SCA2/ATXN2 was detected in a paraffin-embedded section of human colorectal adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-SCA2/ATXN2 Antibody (AAA126189) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)
Figure 2. IHC analysis of SCA2/ATXN2 using anti-SCA2/ATXN2 antibody (AAA126189).SCA2/ATXN2 was detected in a paraffin-embedded section of human thyroiditis tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-SCA2/ATXN2 Antibody (AAA126189) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-ATXN2 antibody	Ataxin-2, the protein encoded by the ATXN2 gene, contains a polyglutamine tract, long expansions (greater than 33 repeats) of which result in spinocerebellar ataxia-2 (SCA2), an autosomal dominant form of olivopontocerebellar atrophy. The gene for spinocerebellar ataxia type 2 (SCA2) has been mapped to 12q24.1. Ataxin-2 associates with L- and T-plastin and that overexpression of ataxin-2 leads to accumulation of T-plastin in mammalian cells.
Product Categories/Family for anti-ATXN2 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	171543895
NCBI GeneID	6311

NCBI Accession #	NP_002964.3
NCBI GenBank Nucleotide #	NM_002973.3
UniProt Accession #	Q99700 ; Q6ZQZ7 ; Q99493 ; A6NLD4
Molecular Weight	140,283 Da
NCBI Official Full Name	ataxin-2
NCBI Official Synonym Full Names	ataxin 2
NCBI Official Symbol	ATXN2
NCBI Official Synonym Symbols	ATX2; SCA2; ASL13; TNRC13
NCBI Protein Information	ataxin-2; trinucleotide repeat containing 13; spinocerebellar ataxia type 2 protein; trinucleotide repeat-containing gene 13 protein
UniProt Protein Name	Ataxin-2
UniProt Gene Name	ATXN2
UniProt Synonym Gene Names	ATX2; SCA2; TNRC13
UniProt Entry Name	ATX2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/sca2-atxn2-antibody-immunohistochemistry-western-blot-elisa/126189>