

Rabbit anti-Human JAM-A/F11R Polyclonal Antibody | anti-F11R antibody

Anti-JAM-A/F11R Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126195	Rabbit	Human	ELISA, Flow Cytometry, Functional Assay, Immunofluorescence, Immunocytochemistry, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

JAM-A/F11R; Polyclonal Antibody; Anti-JAM-A/F11R Antibody Picoband; anti-F11R antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-F11R antibody	ELISA, FCM/FACS (Flow Cytometry), IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human JAM-A/F11R recombinant protein (Position: S28-T294).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




FCM/FACS (Flow Cytometry)

Figure 5. Flow Cytometry analysis of Daudi cells using anti-JAM-A/F11R antibody (AAA126195).Overlay histogram showing Daudi cells stained with AAA126195 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-JAM-A/F11R Antibody (AAA126195, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




IHC (Immunohistochemisry)

Figure 3. IHC analysis of JAM-A/F11R using anti-JAM-A/F11R antibody (AAA126195).JAM-A/F11R was detected in a paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-JAM-A/F11R Antibody (AAA126195) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.




IF (Immunofluorescence)

Figure 4. IF analysis of JAM-A/F11R using anti-JAM-A/F11R antibody (AAA126195).JAM-A/F11R was detected in an immunocytochemical section of T-47D cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-JAM-A/F11R Antibody (AAA126195) overnight at 4 degree C. DyLight488 Conjugated Goat Anti-Rabbit IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



IHC (Immunohiostchemistry)

Figure 2. IHC analysis of JAM-A/F11R using anti-JAM-A/F11R antibody (AAA126195).JAM-A/F11R was detected in a paraffin-embedded section of human colonic adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-JAM-A/F11R Antibody (AAA126195) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of JAM-A/F11R using anti-JAM-A/F11R antibody (AAA126195). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human Hacat whole cell lysates, Lane 2: human T-47D whole cell lysates, Lane 3: human A431 whole cell lysates, Lane 4: human RT4 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-JAM-A/F11R antigen affinity purified polyclonal antibody (#AAA126195) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for JAM-A/F11R at approximately 38 kDa. The expected band size for JAM-A/F11R is at 33 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-F11R antibody

Junctional adhesion molecule A is a protein that in humans is encoded by the F11R gene. Tight junctions represent one mode of cell-to-cell adhesion in epithelial or endothelial cell sheets, forming continuous seals around cells and serving as a physical barrier to prevent solutes and water from passing freely through the paracellular space. The protein encoded by this immunoglobulin superfamily gene member is an important regulator of tight junction assembly in epithelia. In addition, the encoded protein can act as (1) a receptor for reovirus, (2) a ligand for the integrin LFA1, involved in leukocyte transmigration, and (3) a platelet receptor. Multiple 5' alternatively spliced variants, encoding the same protein, have been identified but their biological validity has not been established.

Product Categories/Family for anti-F11R antibody

Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	8393638
NCBI GeneID	50848
NCBI Accession #	NP_058642.1
NCBI GenBank Nucleotide #	NM_016946.4
UniProt Accession #	Q9Y624
Molecular Weight	32,583 Da
NCBI Official Full Name	junctional adhesion molecule A
NCBI Official Synonym Full Names	F11 receptor
NCBI Official Symbol	F11R
NCBI Official Synonym Symbols	JAM; KAT; JAM1; JAMA; JCAM; CD321; PAM-1
NCBI Protein Information	junctional adhesion molecule A; platelet F11 receptor; platelet adhesion molecule 1; junctional adhesion molecule 1

UniProt Protein Name	Junctional adhesion molecule A
UniProt Gene Name	F11R
UniProt Synonym Gene Names	JAM1; JCAM; JAM-A; JAM-1; PAM-1
UniProt Entry Name	JAM1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/jam-a-f11r-antibody-flow-cytometry-immunofluorescence-immunocytochemistry-immunohistochemistry-western-blot-elisa/126195>