

Rabbit anti-Human, Rat EEF2K Polyclonal Antibody | anti-EEF2K antibody

Anti-EEF2K Antibody Picoband

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA126211	Rabbit	Human, Rat	Western Blot, Immunocytochemistry, Immunofluorescence, Flow Cytometry, Functional Assay, ELISA


PRODUCT SYNONYMS

EEF2K; Polyclonal Antibody; Anti-EEF2K Antibody Picoband; anti-EEF2K antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-EEF2K antibody	WB (Western Blot), ICC (Immunocytochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry), ELISA
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human EEF2K recombinant protein (Position: K162-A719).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.

IMAGES




FCM/FACS (Flow Cytometry)

Figure 3. Flow Cytometry analysis of SiHa cells using anti-EEF2K antibody (AAA126211). Overlay histogram showing SiHa cells stained with AAA126211 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-EEF2K Antibody (AAA126211, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IF (Immunofluorescence)

Figure 2. IF analysis of EEF2K using anti-EEF2K antibody (AAA126211). EEF2K was detected in an immunocytochemical section of HepG2 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-EEF2K Antibody (AAA126211) overnight at 4 degree C. DyLight488 Conjugated Goat Anti-Rabbit IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



WB (Western Blot)

Figure 1. Western blot analysis of EEF2K using anti-EEF2K antibody (AAA126211). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human PC-3 whole cell lysates, Lane 2: human U-87MG whole cell lysates, Lane 3: human HepG2 whole cell lysates, Lane 4: rat C6 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-EEF2K antigen affinity purified polyclonal antibody (#AAA126211) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for EEF2K at approximately 82 kDa. The expected band size for EEF2K is at 82 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-EEF2K antibody	Eukaryotic elongation factor-2 kinase (eEF-2 kinase or eEF-2K), also known as calmodulin-dependent protein kinase III (CAMKIII) and calcium/calmodulin-dependent eukaryotic elongation factor 2 kinase, is an enzyme that in humans is encoded by the EEF2K gene. This gene encodes a highly conserved protein kinase in the calmodulin-mediated signaling pathway that links activation of cell surface receptors to cell division. This kinase is involved in the regulation of protein synthesis. It phosphorylates eukaryotic elongation factor 2 (EEF2) and thus inhibits the EEF2 function. The activity of this kinase is increased in many cancers and may be a valid target for anti-cancer treatment.
Product Categories/Family for anti-EEF2K antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	9558749
NCBI GeneID	29904
NCBI Accession #	NP_037434.1
NCBI GenBank Nucleotide #	NM_013302.3
UniProt Accession #	O00418; Q8N588
Molecular Weight	82,144 Da
NCBI Official Full Name	eukaryotic elongation factor 2 kinase
NCBI Official Synonym Full Names	eukaryotic elongation factor-2 kinase
NCBI Official Symbol	EEF2K
NCBI Official Synonym Symbols	eEF-2K; HSU93850

NCBI Protein Information	eukaryotic elongation factor 2 kinase; eEF-2 kinase; elongation factor-2 kinase; eukaryotic elongation factor 2 kinase; calmodulin-dependent protein kinase III; calcium/calmodulin-dependent eukaryotic elongation factor 2 kinase; calcium/calmodulin-depends
UniProt Protein Name	Eukaryotic elongation factor 2 kinase
UniProt Gene Name	EEF2K
UniProt Synonym Gene Names	eEF-2 kinase; eEF-2K
UniProt Entry Name	EF2K_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/eef2k-antibody-western-blot-immunocytochemistry-immunofluorescence-flow-cytometry-elisa/126211>